

DEFORMATION THEORY OF ABELIAN CATEGORIES

WENDY T. LOWEN AND MICHEL VAN DEN BERGH

ABSTRACT. In this paper we develop the basic infinitesimal deformation theory of *abelian categories*. This theory yields a natural generalization of the well-known deformation theory of algebras developed by Gerstenhaber. As part of our deformation theory we define a notion of flatness for abelian categories. We show that various basic properties are preserved under flat deformations and we construct several equivalences between deformation problems.

CONTENTS

1. Introduction	2
2. Preliminaries	4
2.1. Universes	4
2.2. Some constructions depending on the universe	5
2.3. R -linear abelian categories	6
2.4. Derived functors and ind-objects	8
3. Flatness for R -linear abelian categories	9
4. Base change	12
5. Deformations of abelian categories	15
6. Preservation of properties under nilpotent deformations	16
6.1. Nilpotent deformations	17
6.2. Preservation of size	17
6.3. Lifting of objects	18
6.4. Deformations of categories with enough injectives	20
6.5. Deformations of Grothendieck categories	20
6.6. Deformations of locally coherent Grothendieck categories	24
7. Deformation and localization	26
7.1. Statement of the main results	26
7.2. Proofs	27
8. Equivalent deformation problems	30
8.1. Deformation pseudo functors	30
8.2. Abelian and linear deformations	31
8.3. Small skeletons	31
8.4. Elementary deformation equivalences	32
8.5. Ind-objects versus coherent objects	33
8.6. Lifting deformations of localizations	33
8.7. Deformations of Grothendieck categories with acyclic generators	36
8.8. Deformations of categories with enough injectives	37
8.9. Sheaves of modules over a ringed space	37

The first author is an aspirant at the FWO.

The second author is a senior researcher at the FWO.

Appendix A.	38
A.1. Flat nilpotent linear deformations	38
A.2. Deformations with enough injectives	40
Appendix B.	41
References	43

1. INTRODUCTION

In this paper we develop the basic infinitesimal deformation theory of *abelian categories*. This theory yields a natural generalization of the well-known deformation theory of algebras developed by Gerstenhaber [7, 9, 10]. In a subsequent paper [17] we will develop the corresponding obstruction theory in terms of a suitable notion of Hochschild cohomology.

Deformation theory of abelian categories is important for non-commutative algebraic geometry. One of the possible goals of non-commutative algebraic geometry is to understand the abelian (or triangulated) categories which have properties close to those of the (derived) category of (quasi-)coherent sheaves on a scheme. One is particularly interested in those properties which are preserved under suitable deformations. The deformation theory of (abstract) triangulated categories seems at this point somewhat elusive (due to the unclear status of the currently accepted axioms) but, as we will show in this paper, there is a perfectly good deformation theory for abelian categories.

As in any deformation theory we need some kind of flatness in order to control the deformed objects. Therefore the first contribution of this paper is a notion of flatness for abelian categories (see §3). To the best of our knowledge this definition is new.

In the rest of this introduction R is a commutative coherent ring. We will consider R -linear abelian categories. Informally these may be viewed as non-commutative schemes over the (commutative) affine base scheme $\text{Spec } R$.

Let \mathcal{C} be an R -linear abelian category. Our notion of flatness has the following properties:

- (1) if A is an R -algebra then $\text{Mod}(A)$ is flat if and only if A is flat over R ;
- (2) \mathcal{C} is flat if and only if \mathcal{C}^{op} is flat;
- (3) if \mathcal{C} has enough injectives, \mathcal{C} is flat if and only if injectives are flat [3] in \mathcal{C}^{op} ;
- (4) if \mathcal{C} is essentially small, \mathcal{C} is flat if and only if $\text{Lnd}(\mathcal{C})$ is flat (recall that $\text{Lnd}(\mathcal{C})$ is the formal closure of \mathcal{C} under filtered colimits (see 2.2), it is a category with enough injectives);
- (5) flatness is stable under “base change” (see below).

By enlarging the universe (see §2.1) we may assume that any category is small. Therefore, in principle, we could take properties (3) and (4) as the definition of flatness. However this would make the self duality (property (2)) very obscure. Our definition of flatness is somewhat more technical (see §3) but it is manifestly left right symmetric.

Of fundamental importance in algebraic geometry is the concept of base change. There is a natural substitute for this notion in our setting. Consider a morphism $\theta : R \rightarrow S$ between coherent rings such that S is finitely presented over R and

let \mathcal{C} be an R -linear category. The category \mathcal{C}_S is the category of S -objects in \mathcal{C} , i.e. the pairs (C, φ) where $C \in \text{Ob}(\mathcal{C})$ and $\varphi : S \rightarrow \mathcal{C}(C, C)$ is an R -algebra map. Intuitively \mathcal{C}_S the base extension of the “non-commutative scheme” \mathcal{C} to $\text{Spec } S$. We show in §4 that base change is compatible with various natural constructions such as Ind and Mod .

Assume now that θ is surjective such that $I \stackrel{\text{def}}{=} \ker \theta$ satisfies $I^n = 0$ for some n . The surjectivity of θ implies that \mathcal{C}_S is a full abelian subcategory of \mathcal{C} .

Let \mathcal{D} be a flat S -linear category. A flat R -deformation of \mathcal{D} is roughly speaking a flat lift of \mathcal{D} along the functor $(-)_S$. In §6 we show that some of the basic properties of abelian categories are preserved under flat deformation. More precisely, we show that the following properties of \mathcal{D} lift to a flat deformation.

- (1) \mathcal{D} is essentially small;
- (2) \mathcal{D} has enough injectives;
- (3) \mathcal{D} is a Grothendieck category (i.e. a cocomplete abelian category with a generator and exact filtered colimits);
- (4) \mathcal{D} is a locally coherent Grothendieck category (i.e. \mathcal{D} is Grothendieck and is generated by a small abelian subcategory of finitely presented objects).

In addition we show (see Theorem 8.5) that up to equivalence the number of flat deformations of an essentially small, respectively a Grothendieck category is small.

Flatness is necessary for some of these properties. For example if k is a field then there are non-flat deformations of $\text{Mod}(k)$ which do not have enough injectives (Example 6.17) and furthermore the number of non-flat deformations is not small (Remark 8.6).

In §7 we discuss the compatibility of localization with deformations. Among other things we show that a deformation of \mathcal{D} gives rise to deformations of all its localizations (Theorem 7.1).

As a preparation to the sequel of this paper in which we will develop the obstruction theory of abelian categories, we study the associated deformation functors in §8. By $\text{Def}_{\mathcal{D}}(R)$ we denote the flat R -deformations of \mathcal{D} .

We have the following results.

- (1) If \mathcal{D} is an essentially small flat S -linear category then there is an equivalence between $\text{Def}_{\mathcal{D}}(R)$ and $\text{Def}_{\text{Ind}(\mathcal{D})}(R)$.
- (2) If \mathcal{D} is a locally coherent flat S -linear Grothendieck category then there is an equivalence between $\text{Def}_{\mathcal{D}}(R)$ and $\text{Def}_{\text{Fp}(\mathcal{D})}(R)$ where $\text{Fp}(\mathcal{D})$ is the full (abelian) subcategory of \mathcal{D} consisting of the finitely presented objects.

In order to describe some more results of Section §8 we also need to introduce deformations of general R -linear categories. This is done by considering such categories as “rings with several objects” [19]. We denote by $\text{def}_{\mathfrak{b}}(R)$ the flat R -deformations of \mathfrak{b} as R -linear category. Note that deforming an R -linear abelian category \mathcal{D} as *abelian category* is completely different from deforming \mathcal{D} as R -linear category.

We prove the following results.

- (3) If \mathfrak{b} is an essentially small flat S -linear category then there is an equivalence between $\text{def}_{\mathfrak{b}}(R)$ and $\text{Def}_{\text{Mod}(\mathfrak{b})}(R)$ where $\text{Mod}(\mathfrak{b})$ is the category of covariant additive functors from \mathfrak{b} to Ab . In particular if B is a flat S -algebra then there is an equivalence between the deformations of B and the deformations of $\text{Mod}(B)$.

- (4) If \mathcal{D} is a flat S -linear category with enough injectives then there is an equivalence between $\text{Def}_{\mathcal{D}}(R)$ and $\text{def}_{\text{Inj}(\mathcal{D})}(R)$ where $\text{Inj}(\mathcal{D})$ denotes the full (additive) subcategory of \mathcal{D} of injective objects.

Property (3) shows that indeed our deformation theory generalizes the deformation theory of algebras.

In the final section of this paper we apply our methods to the deformations of the category $\text{Mod}(\mathcal{O}_X)$ of sheaves of modules over a ringed space (X, \mathcal{O}_X) . For simplicity of exposition we assume here that \mathcal{O}_X is a sheaf of k -algebras where k is a field. Assume that X has a basis \mathcal{B} satisfying the following acyclicity condition:

$$\forall U \in \mathcal{B} : H^i(U, \mathcal{O}_U) = 0 \quad \text{for } i = 1, 2$$

Let $\mathcal{O}_{\mathcal{B}}$ be the restriction of \mathcal{O}_X to \mathcal{B} and let $\text{PreMod}(\mathcal{O}_{\mathcal{B}})$ be the corresponding category of presheaves. We show that there is an equivalence

$$\text{Def}_{\text{PreMod}(\mathcal{O}_{\mathcal{B}})}(R) \cong \text{Def}_{\text{Mod}(\mathcal{O}_X)}(R)$$

Let \mathfrak{u} be the pre-additive category spanned by the (presheaf) extensions by zero of the \mathcal{O}_U for $U \in \mathcal{B}$. In other words we may take $\text{Ob}(\mathfrak{u}) = \mathcal{B}$ and we have

$$\mathfrak{u}(U, V) = \begin{cases} \mathcal{O}_X(U) & \text{if } U \subset V \\ 0 & \text{otherwise} \end{cases}$$

Using property (3) above, it is easy to see that

$$\text{Def}_{\text{PreMod}(\mathcal{O}_{\mathcal{B}})}(R) \cong \text{def}_{\mathfrak{u}}(R)$$

These results confirm the fundamental insight of Gerstenhaber and Schack [6, 8] that one should define the deformations of a ringed space (X, \mathcal{O}_X) not as the deformations of \mathcal{O}_X as a sheaf of k -algebras, but rather as the deformations of the k -linear category \mathfrak{u} (or of the “diagram” $(\mathcal{B}, \mathcal{O}_{\mathcal{B}})$ in case $X \in \mathcal{B}$). These “virtual” deformations are nothing but the deformations of the abelian category $\text{Mod}(\mathcal{O}_X)$.

2. PRELIMINARIES

2.1. Universes. It is well-known [18] that category theory needs some extension of the Zermelo-Fraenkel axioms of set theory (ZF). One possible extension is given by the Gödel-Bernays axioms (GB) which incorporates classes into set theory. This makes it possible to introduce the category Set while at the same time avoiding Russel’s paradox.

This solution is not entirely satisfying since for example one would also like to talk about Cls , the category of all classes, and there is no room for this notion in GB. In particular, in the deformation theory of categories we consider below, this seems to lead to foundational problems.

To solve such problems Grothendieck introduced a more flexible extension of the Zermelo-Fraenkel system: the theory of universes [2]. The theory of universes does not introduce new types of objects but adds the universe axiom (U) below.

A universe \mathcal{U} is a set with the following properties:

- (1) if $x \in \mathcal{U}$ and if $y \in x$ then $y \in \mathcal{U}$;
- (2) if $x, y \in \mathcal{U}$ then $\{x, y\} \in \mathcal{U}$;
- (3) if $x \in \mathcal{U}$ then the powerset $\mathcal{P}(x)$ of x is in \mathcal{U} ;
- (4) if $(x_i)_{i \in I}$ is a family of objects in \mathcal{U} indexed by an element of \mathcal{U} then $\bigcup_{i \in I} x_i \in \mathcal{U}$;
- (5) if $U \in \mathcal{U}$ and $f : U \rightarrow \mathcal{U}$ is a function, then $\{f(x) \mid x \in U\} \in \mathcal{U}$.

Note that (x, y) is defined as $\{\{x, y\}, x\}$ and hence if $x, y \in \mathcal{U}$ then so is (x, y) . A universe \mathcal{U} with $\mathbb{N} \in \mathcal{U}$ is itself a model for ZF.

As the only known non-empty universe only contains finite sets we need the following new axiom:

(U) every set is the element of a universe.

In this paper, we will work with ZFCU (the ZF axioms + the axiom of choice + the universe axiom). By requiring $\{x, \mathbb{N}\} \in \mathcal{U}$, every set x is the element of a universe containing x and \mathbb{N} . From now on, by a universe we will always mean a universe containing \mathbb{N} . In particular, every such universe is itself a model for ZFC.

We now recall some terminology.

- Definition 2.1.**
- (1) A set or cardinal is \mathcal{U} -small if it has the same cardinality as an element of \mathcal{U} .
 - (2) A category \mathcal{C} consists of a set(!) of “objects” $\text{Ob}(\mathcal{C})$ and a set of “arrows” $\text{Mor}(\mathcal{C})$ with the usual extra structure.
 - (3) \mathcal{U} -Set is the category whose objects consist of elements of \mathcal{U} and whose Hom-sets are just the standard Hom’s between sets. Likewise if \mathcal{E} is a “structure” [5] (e.g. abelian groups or rings) then \mathcal{U} - \mathcal{E} is the category of \mathcal{E} -objects whose underlying set is in \mathcal{U} . In particular, \mathcal{U} -Cat (resp. \mathcal{U} -Gd) is the category of categories (resp. groupoids) \mathcal{C} with $\text{Ob}(\mathcal{C}) \in \mathcal{U}$ and $\text{Mor}(\mathcal{C}) \in \mathcal{U}$ and the usual Hom’s between categories.
 - (4) A category is \mathcal{U} -small if both its objects and arrows are \mathcal{U} -small sets.
 - (5) A category is *essentially \mathcal{U} -small* if it equivalent to a \mathcal{U} -small category.
 - (6) A category is a \mathcal{U} -category if it has \mathcal{U} -small Hom-sets.
 - (7) An abelian \mathcal{U} -category \mathcal{C} is *\mathcal{U} -Grothendieck* if \mathcal{C} has a generator; \mathcal{U} -small colimits exist in \mathcal{C} and \mathcal{U} -small filtered colimits are exact.

Remark 2.2. If $E \in \text{Ob}(\mathcal{U}\text{-}\mathcal{E})$, then $E \in \mathcal{U}$ since E is described by an element of \mathcal{U} . For example, $\mathcal{C} \in \text{Ob}(\mathcal{U}\text{-Cat})$ is described by an element of $\mathcal{U}^{\times 6} \subset \mathcal{U}$.

Lemma 2.3. For $\mathcal{C}, \mathcal{D} \in \mathcal{U}\text{-Cat}$, $\text{Hom}(\mathcal{C}, \mathcal{D}) \in \mathcal{U}\text{-Cat}$. □

Remark 2.4. The axiom of choice allows us to replace a \mathcal{U} -category \mathcal{C} with an isomorphic category \mathcal{C}' with $\text{Ob}(\mathcal{C}') = \text{Ob}(\mathcal{C})$ and $\mathcal{C}'(C, D) \in \mathcal{U}$ for every $C, D \in \text{Ob}(\mathcal{C})$. In particular, if \mathcal{C} is a (pre-additive) \mathcal{U} -category, we can define representable functors

$$\mathcal{C}(C, -) : \mathcal{C} \longrightarrow \mathcal{U}\text{-Set}(\mathcal{U}\text{-Ab}) : D \longmapsto \mathcal{C}'(C, D)$$

where \mathcal{C}' is as above.

The universe axiom is the basis for the very useful “extension of the universe” principle. I.e. by selecting a large enough universe we may assume that any individual category is small. The theory of universes comes at a price however, namely the dependence of the notations on the chosen universe. Since this is rather tedious one usually fixes the universe in advance and then drops it from the notations except when invoking the extension of the universe principle. We will follow these conventions in this paper.

2.2. Some constructions depending on the universe. If \mathfrak{a} is a pre-additive category then we denote by $\mathcal{U}\text{-Mod}(\mathfrak{a})$ the category of covariant additive functors from \mathfrak{a} to $\mathcal{U}\text{-Ab}$. It is easy to see that if \mathfrak{a} is essentially \mathcal{U} -small then $\mathcal{U}\text{-Mod}(\mathfrak{a})$ is a \mathcal{U} -Grothendieck category. If \mathfrak{a} is a \mathcal{U} -category, $\mathcal{U}\text{-Mod}(\mathfrak{a})$ contains functors

$\mathfrak{a}(A, -) : \mathfrak{a} \rightarrow \mathcal{U}\text{-Ab}$ for $A \in \mathfrak{a}$ (see Remark 2.4). In this case we define $\mathcal{U}\text{-mod}(\mathfrak{a})$ as the full subcategory of $\mathcal{U}\text{-Mod}(\mathfrak{a})$ containing all functors that can be written as cokernels of maps $\bigoplus_{i=1}^m \mathfrak{a}(A_i, -) \rightarrow \bigoplus_{j=1}^n \mathfrak{a}(B_j, -)$. If $\mathcal{U} \subset \mathcal{V}$ is an inclusion of universes, then Yoneda's lemma yields an equivalence of categories $\mathcal{U}\text{-mod}(\mathfrak{a}) \rightarrow \mathcal{V}\text{-mod}(\mathfrak{a})$.

If \mathcal{C} is an essentially small \mathcal{U} -category then $\mathcal{U}\text{-Ind}(\mathcal{C})$ is the full subcategory of $\mathcal{U}\text{-Mod}(\mathcal{C})$ consisting of left exact functors. It is well-known that $\mathcal{U}\text{-Ind}(\mathcal{C})$ is a \mathcal{U} -Grothendieck category. The objects in $\mathcal{U}\text{-Ind}(\mathcal{C})$ may be written as formal \mathcal{U} -small filtered colimits of objects in \mathcal{C} and the Hom-sets are computed by the rule

$$\text{Hom}_{\mathcal{U}\text{-Ind}(\mathcal{C})}(\text{colim}_{i \in I} A_i, \text{colim}_{j \in J} B_j) = \lim_{i \in I} \text{colim}_{j \in J} \text{Hom}_{\mathcal{C}}(A_i, B_j).$$

An object C in a \mathcal{U} -category \mathcal{C} is \mathcal{U} -finitely presented if the functor $\mathcal{C}(C, -) : \mathcal{C} \rightarrow \mathcal{U}\text{-Set}$ (see Remark 2.4) preserves \mathcal{U} -small filtered colimits. We define $\mathcal{U}\text{-Fp}(\mathcal{C})$ as the full subcategory of \mathcal{C} containing precisely the \mathcal{U} -finitely presented objects. It is well-known that if \mathcal{C} contains a \mathcal{U} -small full subcategory \mathfrak{g} of \mathcal{U} -finitely presented generators of \mathcal{C} , then $\mathcal{U}\text{-Fp}(\mathcal{C})$ is the finite colimit closure of \mathfrak{g} in \mathcal{C} [4] and in particular is essentially \mathcal{U} -small. If \mathfrak{a} is essentially \mathcal{U} -small, it is well-known and easy to see that $\mathcal{U}\text{-Fp}(\mathcal{U}\text{-Mod}(\mathfrak{a})) = \mathcal{U}\text{-mod}(\mathfrak{a})$.

A \mathcal{U} -Grothendieck category \mathcal{C} is locally coherent if it has a \mathcal{U} -small set of \mathcal{U} -finitely presented generators and $\mathcal{U}\text{-Fp}(\mathcal{C})$ is a (necessarily essentially \mathcal{U} -small) abelian category.

If \mathcal{C} is a locally coherent \mathcal{U} -Grothendieck category then the natural functor

$$\mathcal{U}\text{-Ind}(\mathcal{U}\text{-Fp}(\mathcal{C})) \rightarrow \mathcal{C}$$

is an equivalence of categories. If \mathcal{C} is essentially \mathcal{U} -small then the natural functor

$$\mathcal{C} \rightarrow \mathcal{U}\text{-Fp}(\mathcal{U}\text{-Ind}(\mathcal{C}))$$

is an equivalence as well.

Convention. From now on we work with a fixed universe \mathcal{U} . All categories will be \mathcal{U} -categories. The notions of small and essentially small are with respect to \mathcal{U} . The same holds for the notion of a Grothendieck category. The symbols Mod , Ind , Fp and mod are implicitly prefixed by \mathcal{U} . Individual objects such as abelian groups, rings, modules are, unless otherwise specified, assumed to be \mathcal{U} -small.

2.3. R -linear abelian categories. Consider a commutative ring R . Recall that an R -linear category is a pre-additive category \mathfrak{a} together with a ring map $\rho : R \rightarrow \text{Nat}(1_{\mathfrak{a}}, 1_{\mathfrak{a}})$. ρ induces a ring map $\rho_A : R \rightarrow \mathfrak{a}(A, A)$ for every object A and an action of R on every Hom-set. This leads to the equivalent definition of an R -linear category as a category enriched in the category $\text{Mod}(R)$ of R -modules. A pre-additive category is of course the same as a \mathbb{Z} -linear category.

Unless otherwise stated, we will assume the ring R to be coherent. Let $\text{mod}(R)$ denote the full abelian subcategory of $\text{Mod}(R)$ of finitely presented R -modules. Consider an abelian R -linear category \mathcal{C} . For every object C of \mathcal{C} we obtain a (up to a canonical natural isomorphism) unique finite colimit preserving functor

$$(2.1) \quad (- \otimes_R C) : \text{mod}(R) \rightarrow \mathcal{C}$$

with $R \otimes_R C = C$. We can construct its left derived functors $\text{Tor}_i^R(-, C)$ using projective resolutions in $\text{mod}(R)$. In fact, we naturally have (R -bilinear) bifunctors

$$\text{Tor}_i^R(-, -) : \text{mod}(R) \times \mathcal{C} \rightarrow \mathcal{C}.$$

The functors $(X \otimes_R -)$ are finite colimit preserving. It is easily seen (using a fixed free resolution of X) that the functors $\mathrm{Tor}_i^R(X, -)$ form a homological δ -functor.

In a completely analogous way we define

$$\mathrm{Hom}_R(-, C) : (\mathrm{mod}(R))^{\mathrm{op}} \longrightarrow \mathcal{C}$$

as the unique finite limit preserving functor with $\mathrm{Hom}_R(R, C) = C$, and taking its right derived functors (again using projective resolutions in $\mathrm{mod}(R)$) we obtain (R -bilinear) bifunctors

$$\mathrm{Ext}_R^i(-, -) : (\mathrm{mod}(R))^{\mathrm{op}} \times \mathcal{C} \longrightarrow \mathcal{C}.$$

The functors $\mathrm{Hom}_R(X, -)$ are finite limit preserving. The functors $\mathrm{Ext}_R^i(X, -)$ form a cohomological δ -functor.

Note that $\mathrm{Hom}_R(X, -) : \mathcal{C} \longrightarrow \mathcal{C}$ is in fact nothing but the opposite of the tensor functor $(X \otimes_R -) : \mathcal{C}^{\mathrm{op}} \longrightarrow \mathcal{C}^{\mathrm{op}}$ of $\mathcal{C}^{\mathrm{op}}$.

Definition 2.5. [3] An object C of \mathcal{C} is *flat (over R)* if the functor $(- \otimes_R C) : \mathrm{mod}(R) \longrightarrow \mathcal{C}$ is exact. Dually, C is *coflat (over R)* if $\mathrm{Hom}_R(-, C) : \mathrm{mod}(R) \longrightarrow \mathcal{C}$ is exact.

Remark 2.6. Flatness of C is equivalent to the vanishing of $\mathrm{Tor}_1^R(-, C)$ and to the vanishing of all $\mathrm{Tor}_i^R(-, C)$ for $i \geq 1$. Dually, coflatness of C is equivalent to the vanishing of $\mathrm{Ext}_R^1(-, C)$ and to the vanishing of all $\mathrm{Ext}_R^i(-, C)$ for $i \geq 1$. Consequently, in an exact sequence $0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$, if B and C are flat, the same holds for A , and dually, if A and B are coflat, the same holds for C .

We will now list some useful facts concerning the Hom_R and \otimes_R functors.

Proposition 2.7. Consider an R -linear functor $F : \mathcal{C} \longrightarrow \mathcal{D}$ between abelian R -linear categories. If F is exact, then for $X \in \mathrm{mod}(R)$ and $C \in \mathcal{C}$

$$(2.2) \quad \mathrm{Ext}_R^i(X, F(C)) = F(\mathrm{Ext}_R^i(X, C)).$$

If F is only left exact, then (2.2) holds for $i = 0$.

Proof. This follows if we replace X by a free resolution. \square

Corollary 2.8. Consider an R -linear category \mathcal{C} and a small category \mathcal{I} . If \mathcal{I} -colimits are exact in \mathcal{C} , $\mathrm{Ext}_R^i(X, -)$ preserves them for $X \in \mathrm{mod}(R)$.

Proof. By taking F to be the functor $\mathrm{colim} : \mathrm{Fun}(\mathcal{I}, \mathcal{C}) \longrightarrow \mathcal{C}$, this follows from Proposition 2.7. \square

Proposition 2.9. For $X, Y \in \mathrm{mod}(R)$ and $C, D, E \in \mathcal{C}$, the following hold:

- (0) $\mathcal{C}(X \otimes_R C, D) \cong \mathrm{Hom}_R(X, \mathcal{C}(C, D)) \cong \mathcal{C}(C, \mathrm{Hom}_R(X, D))$;
- (1) $\mathrm{Hom}_R(X \otimes_R Y, C) \cong \mathrm{Hom}_R(X, \mathrm{Hom}_R(Y, C))$;
- (2) $(X \otimes_R Y) \otimes_R C \cong X \otimes_R (Y \otimes_R C)$;
- (3) if C is coflat, $X \otimes_R \mathrm{Hom}_R(Y, C) \cong \mathrm{Hom}_R(\mathrm{Hom}_R(X, Y), C)$;
- (4) if D is flat, $\mathrm{Hom}_R(X, Y \otimes_R D) \cong \mathrm{Hom}_R(X, Y) \otimes_R D$;
- (5) if E is injective, $X \otimes_R \mathcal{C}(C, E) \cong \mathcal{C}(\mathrm{Hom}_R(X, C), E)$;
- (6) if E is injective and C is coflat, then $\mathcal{C}(C, E)$ is flat;
- (7) if E is injective and C is flat, then $\mathcal{C}(C, E)$ is coflat;
- (8) if $\mathrm{Ext}_R^1(C, Z \otimes_R E) = 0$ for all Z in $\mathrm{mod}(R)$ and E is flat, then $X \otimes_R \mathcal{C}(C, E) = \mathcal{C}(C, X \otimes_R E)$ and $\mathcal{C}(C, E)$ is flat. \square

Consider a finitely generated ideal I of R . For every $C \in \mathcal{C}$, we obtain a map $I \otimes_R C \rightarrow C$, which is not necessarily mono unless C is flat. We will denote the image of this map by IC . Dually, we will denote the image of the map $C \rightarrow \text{Hom}_R(I, C)$ by CI . It is easy to see that $(IJ)C = I(JC)$ as subobjects of C .

If (\mathfrak{a}, ρ) is an R -linear category and $F \in \text{Mod}(\mathfrak{a})$, then the abelian groups $F(A)$ inherit an R -module structure from the maps $\rho(r) : A \rightarrow A$. In fact, $\text{Mod}(\mathfrak{a})$ is isomorphic to the R -linear category of R -linear functors from \mathfrak{a} into $\text{Mod}(R)$ (see also §4).

Proposition 2.10. *For an essentially small R -linear category \mathfrak{a} , the Hom_R and \otimes_R functors on $\text{Mod}(\mathfrak{a})$ are computed pointwise. I.e.*

$$\begin{aligned} (X \otimes_R F)(A) &= X \otimes_R F(A) \\ \text{Hom}_R(X, F)(A) &= \text{Hom}_R(X, F(A)) \end{aligned}$$

for $X \in \text{mod}(R)$, $F \in \text{Mod}(\mathfrak{a})$ and $A \in \mathfrak{a}$. In particular, F in $\text{Mod}(\mathfrak{a})$ is flat (resp. coflat) if and only if all its values $F(A)$ are flat (resp. coflat) in $\text{Mod}(R)$. \square

2.4. Derived functors and ind-objects. Some arguments in this paper are based on extending derived functors to ind-objects (see §2.2). Therefore in this section we discuss some of the relevant properties in this regard.

Definition 2.11. (1) A functor $F : \mathcal{A} \rightarrow \mathcal{B}$ between an arbitrary category \mathcal{A} and a pre-additive category \mathcal{B} is called *effaceable*[11] if for every $A \in \mathcal{A}$ there is a monomorphism $u : A \rightarrow A'$ with $F(u) = 0$. F is called *co-effaceable* if for every A there is an epimorphism $u : A' \rightarrow A$ with $F(u) = 0$.
(2) A functor $F : \mathcal{A} \rightarrow \mathbf{Ab}$ is called *weakly effaceable* if for each $A \in \mathcal{A}$ and $x \in F(A)$ there is a monomorphism $u : A \rightarrow A'$ with $F(u)(x) = 0$.

Proposition 2.12. [11] *Let $F : \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor between abelian categories. If \mathcal{A} has enough injectives then the following are equivalent:*

- (1) F is effaceable;
- (2) $F(I) = 0$ for all injectives I of \mathcal{A} . \square

Consider an extension to ind-objects

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & \text{Ind}(\mathcal{C}) \\ F \downarrow & & \downarrow \text{Ind}(F) \\ \mathcal{D} & \longrightarrow & \text{Ind}(\mathcal{D}) \end{array}$$

where \mathcal{C}, \mathcal{D} are essentially small abelian categories.

Proposition 2.13. *The following are equivalent:*

- (1) F is effaceable;
- (2) $\text{Ind}(F)$ is effaceable.

Proof. Suppose $\text{Ind}(F)$ is effaceable and consider C in \mathcal{C} . There is a monomorphism $u : C \rightarrow \text{colim}_i C_i$ in $\text{Ind}(\mathcal{C})$ with $\text{Ind}(F)(u) = 0$ for certain C_i in \mathcal{C} . Consider the maps $s_j : C_j \rightarrow \text{colim}_i C_i$. Since C is finitely presented, u factorizes as $u = s_j \circ f$ for some $f : C \rightarrow C_j$. Now s_j is mapped by $\text{Ind}(F)$ onto the canonical map $F(C_j) \rightarrow \text{colim}_i F(C_i)$, and its composition with $F(f)$ is zero. Since $F(C)$ is finitely presented

in $\text{Ind}(\mathcal{D})$, there is a map $s_{jk} : C_j \rightarrow C_k$ such that $F(s_{jk}) \circ F(f) = 0$, so $s_{jk} \circ f$ is a monomorphism that serves our purpose.

Now suppose F is effaceable. By Proposition 2.12 it suffices to prove that $\text{Ind}(F)(I) = 0$ for every injective $I = \text{colim}_i C_i$ of $\text{Ind}(\mathcal{C})$. For every C_i , we take a monomorphism $u_i : C_i \rightarrow C'_i$ in \mathcal{C} with $F(u_i) = 0$. Since I is injective, the maps $s_j : C_j \rightarrow I$ factorize as $s_j = f_j \circ u_j$. It follows that all the maps $\text{Ind}(F)(s_j) : F(C_j) \rightarrow \text{colim}_i F(C_i)$ are zero and hence $\text{Ind}(F)(I) = \text{colim}_i F(C_i) = 0$. \square

If \mathcal{C} is an abelian category then $\text{Ext}_{\mathcal{C}}^i(-, -)$ denotes the *Yoneda* Ext-groups between \mathcal{C} -objects. If \mathcal{C} has enough injectives, then $\text{Ext}_{\mathcal{C}}^i(C, -)_i$ are the derived functors of $\mathcal{C}(C, -)$. The following proposition is presumably well-known, but we have been unable to find a reference.

Proposition 2.14. *Assume that \mathcal{C} is essentially small. For $A, B \in \mathcal{C}$, we have*

$$(2.3) \quad \text{Ext}_{\mathcal{C}}^i(A, B) = \text{Ext}_{\text{Ind}(\mathcal{C})}^i(A, B).$$

Proof. The formula (2.3) is obviously true for $i = 0$. Then by [12, Lemma II.2.1.3], it is sufficient to show that the functor $\text{Ext}_{\text{Ind}(\mathcal{C})}^i(-, B)$ restricted to \mathcal{C} is weakly effaceable. This follows from Proposition 2.15 below with $G = \text{Ext}_{\text{Ind}(\mathcal{C})}^i(-, B)$. \square

Proposition 2.15. *Assume that \mathcal{C} is essentially small. If in the following commutative diagram*

$$\begin{array}{ccc} \mathcal{C}^{\text{op}} & \longrightarrow & (\text{Ind}(\mathcal{C}))^{\text{op}} \\ & \searrow F & \downarrow G \\ & & \text{Ab} \end{array}$$

G is weakly effaceable, the same holds for F .

Proof. Consider C in \mathcal{C} and $x \in F(C)$. There is an epimorphism $f : \text{colim}_i C_i \rightarrow C$ in $\text{Ind}(\mathcal{C})$ with $G(f)(x) = 0$ for certain C_i in \mathcal{C} . Consider the maps $s_j : C_j \rightarrow \text{colim}_i C_i$. We have that $C = \sum_i \text{Im}(f \circ s_i)$, but since C is finitely generated in $\text{Ind}(\mathcal{C})$, we find an epimorphism $f \circ s_j : C_j \rightarrow C$ in $\text{Ind}(\mathcal{C})$ (and thus in \mathcal{C}) with $F(f \circ s_j)(x) = G(f \circ s_j)(x) = G(s_j)(G(f)(x)) = 0$. \square

3. FLATNESS FOR R -LINEAR ABELIAN CATEGORIES

An R -linear category is called *flat* if its Hom-sets are flat in $\text{Mod}(R)$. In this section, we introduce a different notion of flatness for an R -linear *abelian* category \mathcal{C} . This notion has the following properties:

- (1) \mathcal{C} is flat if and only if \mathcal{C}^{op} is flat;
- (2) if \mathcal{C} has enough injectives, \mathcal{C} is flat if and only if injectives are coflat;
- (3) if \mathcal{C} is essentially small, \mathcal{C} is flat if and only if $\text{Ind}(\mathcal{C})$ is flat;
- (4) if \mathfrak{a} is an essentially small R -linear category, \mathfrak{a} is flat (as an R -linear category) if and only if $\text{Mod}(\mathfrak{a})$ is flat (as an abelian R -linear category).

By enlarging the universe (see §2.1) we may assume that any category is small. Therefore, in principle, we could take properties (3) and (4) as the definition of flatness. However this would make the self duality (property (2)) very obscure. Therefore we give a somewhat more technical, but manifestly left right symmetric definition below.

Let \mathcal{C} be an R -linear abelian category. The following result is crucial for what follows.

Proposition 3.1. *For $X \in \text{mod}(R)$, the following are equivalent:*

- (1) $\text{Ext}_R^1(X, -) : \mathcal{C} \rightarrow \mathcal{C}$ is effaceable;
- (2) $\text{Ext}_R^1(X, \mathcal{C}(-, -)) : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathbf{Ab}$ is weakly effaceable;
- (3) $\text{Tor}_1^R(X, -) : \mathcal{C} \rightarrow \mathcal{C}$ is co-effaceable.

Proof. Since (2) is clearly self-dual, and (3) is the dual statement of (1), it suffices to prove the equivalence of (1) and (2). Take a projective resolution P of X and let Y denote the kernel of $P_0 \rightarrow X$. Note that since

$$\text{Ext}^1(X, \mathcal{C}(A, B)) = H_1\mathcal{C}(A, \text{Hom}_R(P, B)),$$

an element in this group can be represented by a \mathcal{C} -map $A \rightarrow \text{Hom}_R(P_1, B)$ such that the composition with $\text{Hom}_R(P_1, B) \rightarrow \text{Hom}_R(P_2, B)$ is zero. Suppose (2) holds. Consider an arbitrary \mathcal{C} -map $A \rightarrow \text{Ext}_R^1(X, B)$. We will show that there is a mono $B \rightarrow B'$ such that $A \rightarrow \text{Ext}_R^1(X, B) \rightarrow \text{Ext}_R^1(X, B')$ is zero, which proves (1). Consider the pullback

$$\begin{array}{ccc} A' & \longrightarrow & A \\ \downarrow & & \downarrow \\ \text{Hom}_R(Y, B) & \longrightarrow & \text{Ext}_R^1(X, B). \end{array}$$

The composition $A' \rightarrow \text{Hom}_R(Y, B) \rightarrow \text{Hom}_R(P_1, B)$ represents an element of $\text{Ext}^1(X, \mathcal{C}(A, B))$, hence by (2) we can take $A'' \rightarrow A'$ and $B \rightarrow B'$ such that $A'' \rightarrow A' \rightarrow \text{Hom}_R(Y, B) \rightarrow \text{Hom}_R(Y, B')$ factors over $\text{Hom}_R(P_0, B') \rightarrow \text{Hom}_R(Y, B')$. It follows that the composition $A'' \rightarrow A' \rightarrow \text{Hom}_R(Y, B) \rightarrow \text{Hom}_R(Y, B') \rightarrow \text{Ext}_R^1(X, B')$ is zero, and consequently $A \rightarrow \text{Ext}_R^1(X, B) \rightarrow \text{Ext}_R^1(X, B')$ is zero too.

Now suppose (1) holds and consider an element in $\text{Ext}^1(X, \mathcal{C}(A, B))$ represented by $A \rightarrow \text{Hom}_R(P_1, B)$. Take a monomorphism $B \rightarrow B'$ such that $\text{Ext}_R^1(X, B) \rightarrow \text{Ext}_R^1(X, B')$ is zero. Let K denote the kernel of $\text{Hom}_R(P_1, B) \rightarrow \text{Hom}_R(P_2, B)$ and let L' denote the image of $\text{Hom}_R(P_0, B') \rightarrow \text{Hom}_R(P_1, B')$. We obtain the following factorization

$$\begin{array}{ccccc} A' & \longrightarrow & A & & \\ \downarrow & & \downarrow & \searrow & \\ & & K & \longrightarrow & \text{Hom}_R(P_1, B) \\ \downarrow & & \downarrow & & \downarrow \\ \text{Hom}_R(P_0, B') & \longrightarrow & L' & \longrightarrow & \text{Hom}_R(P_1, B'), \end{array}$$

in which A' is defined as a pullback. Clearly, the composition $A' \rightarrow \text{Hom}_R(P_1, B')$ represents zero in $\text{Ext}^1(X, \mathcal{C}(A', B'))$. \square

Definition 3.2. The category \mathcal{C} is called *flat (over R)* if every $X \in \text{mod}(R)$ satisfies the equivalent conditions of Proposition 3.1.

Note that the notion of flatness is independent of the choice of universe.

Proposition 3.3. *The following are equivalent:*

- (1) \mathcal{C} is flat over R ;
- (2) \mathcal{C}^{op} is flat over R .

Proof. Immediate from Proposition 3.1. □

Proposition 3.4. *If \mathcal{C} has enough injectives, the following are equivalent:*

- (1) \mathcal{C} is flat over R ;
- (2) injectives in \mathcal{C} are coflat.

Proof. This follows from Proposition 2.12 with $F = \text{Ext}_R^1(X, -)$. □

Proposition 3.5. *If \mathcal{C} is flat (over R), the following hold:*

- (1) For every $X \in \text{mod}(R)$ and $i \geq 1$, $\text{Ext}_R^i(X, -)$ is effaceable;
- (2) For every $X \in \text{mod}(R)$, the cohomological δ -functor $(\text{Ext}_R^i(X, -))_{i \geq 0}$ is universal [11];
- (3) for every $X \in \text{mod}(R)$ and $i \geq 1$, $\text{Tor}_i^R(X, -)$ is co-effaceable.
- (4) for every $X \in \text{mod}(R)$, the homological δ -functor $(\text{Tor}_i^R(X, -))_{i \geq 0}$ is universal.

Proof. (1) and (3) follow by dimension shifting in the first argument. (2) and (4) follow from (1) and (3) by [11]. □

The category of ind-objects $\text{Ind}(\mathcal{C})$ (see §2.2) is obviously R -linear. It follows from Proposition 2.7 that the functor $\mathcal{C} \rightarrow \text{Ind}(\mathcal{C})$ commutes with Tor^R and Ext_R . So $\text{Tor}_i^{\text{Ind}(\mathcal{C}), R}(X, -)$ and $\text{Ext}_{\text{Ind}(\mathcal{C}), R}^i(X, -)$ are the extensions to ind-objects of $\text{Tor}_i^{\mathcal{C}, R}(X, -)$ and $\text{Ext}_{\mathcal{C}, R}^i(X, -)$.

Proposition 3.6. *Assume that \mathcal{C} is essentially small. The following are equivalent:*

- (1) \mathcal{C} is flat over R ;
- (2) $\text{Ind}(\mathcal{C})$ is flat over R .

Proof. This follows immediately from Proposition 2.13 with $F = \text{Ext}_{\mathcal{C}, R}^1(X, -)$ and $\text{Ind}(F) = \text{Ext}_{\text{Ind}(\mathcal{C}), R}^1(X, -)$. □

Proposition 3.7. *For an essentially small R -linear category \mathfrak{a} , the following are equivalent:*

- (1) \mathfrak{a} is flat over R ;
- (2) the abelian category $\text{Mod}(\mathfrak{a})$ is flat over R .

Proof. First, note that for every E in $\text{Mod}(\mathfrak{a})$ and for every A in \mathfrak{a} ,

$$\begin{aligned} \text{Mod}(R)(X, E(A)) &= \text{Mod}(R)(X, \text{Mod}(\mathfrak{a})(\mathfrak{a}(A, -), E)) \\ &= \text{Mod}(\mathfrak{a})(X \otimes_R \mathfrak{a}(A, -), E). \end{aligned}$$

By Proposition 2.10, $\text{Mod}(\mathfrak{a})$ is flat if and only if for every injective E in $\text{Mod}(\mathfrak{a})$ and every A in \mathfrak{a} , $E(A)$ is coflat. Also by Proposition 2.10, \mathfrak{a} is flat if and only if every functor $\mathfrak{a}(A, -)$ for A in \mathfrak{a} is flat. And since $\text{Mod}(\mathfrak{a})$ has enough injectives, this is equivalent to requiring that for every injective E in $\text{Mod}(\mathfrak{a})$ the functor $\text{Mod}(\mathfrak{a})(-, E) \circ (- \otimes_R \mathfrak{a}(A, -)) : \text{mod}(R) \rightarrow \text{Mod}(R)$ is exact. The statement then follows from the computation above. □

4. BASE CHANGE

In this section we study a natural substitute for the notion of base change in algebraic geometry, and we show that it is compatible with constructions such as Lnd and Mod .

We fix a homomorphism of commutative rings $\theta : R \rightarrow S$. For $M \in \text{Mod}(S)$, \overline{M} denotes M considered as an R -module using θ . For an S -linear category \mathfrak{b} , $\overline{\mathfrak{b}}$ is the R -linear category with $\text{Ob}(\overline{\mathfrak{b}}) = \text{Ob}(\mathfrak{b})$ and $\overline{\mathfrak{b}}(B, B') = \overline{\mathfrak{b}(B, B')}$. More often than not we will just write \mathfrak{b} for $\overline{\mathfrak{b}}$.

For an R -linear category \mathfrak{a} , $S \otimes_R \mathfrak{a}$ is the S -linear category with $\text{Ob}(S \otimes_R \mathfrak{a}) = \text{Ob}(\mathfrak{a})$ and $(S \otimes_R \mathfrak{a})(A, A') = S \otimes_R \mathfrak{a}(A, A')$.

The functor $S \otimes_R (-)$ is left adjoint to $\overline{(-)}$ in the sense that for an R -linear category \mathfrak{a} and an S -linear category \mathfrak{b} , there is an isomorphism, natural in \mathfrak{a} , \mathfrak{b} :

$$\overline{\text{Add}(S)(S \otimes_R \mathfrak{a}, \mathfrak{b})} \cong \text{Add}(T)(\mathfrak{a}, \overline{\mathfrak{b}})$$

as R -linear categories where $\text{Add}(T)$ denotes the T -linear functors.

For an R -linear category (\mathcal{C}, ρ) , let \mathcal{C}_S be the following S -linear category of S -objects: the objects of \mathcal{C}_S are couples (C, φ) where C is an object of \mathcal{C} and $\varphi : S \rightarrow \mathcal{C}(C, C)$ is a ring map with $\varphi \circ \theta = \rho_C$. The morphisms of \mathcal{C}_S are the obvious compatible \mathcal{C} -morphisms. Clearly \mathcal{C}_S becomes S -linear using the ring maps φ . The functor $(-)_S$ is right adjoint to $\overline{(-)}$ in the sense that for an R -linear category \mathcal{C} and an S -linear category \mathcal{D} , there is an isomorphism, natural in \mathcal{C} , \mathcal{D} :

$$\overline{\text{Add}(S)(\mathcal{D}, \mathcal{C}_S)} \cong \text{Add}(R)(\overline{\mathcal{D}}, \mathcal{C})$$

of R -linear categories.

Remark 4.1. It is readily seen that $(\text{Mod}(R))_S \cong \text{Mod}(S)$, hence we deduce the isomorphism $\text{Add}(S)(\mathfrak{b}, \text{Mod}(S)) \cong \text{Mod}(\mathfrak{b})$ for an S -linear category \mathfrak{b} which was already mentioned just before Proposition 2.10.

Proposition 4.2. *The forgetful functor $F : \mathcal{C}_S \rightarrow \mathcal{C}$ preserves, reflects and creates [1] all limits and colimits.*

Proof. We give the construction of limits in \mathcal{C}_S that exist in \mathcal{C} . Let $G : \mathcal{I} \rightarrow \mathcal{C}_S$ with $G(i) = (C_i, \varphi_i)$ be an arbitrary functor and let $\lim_i C_i$ denote the limit of $F \circ G$. For every $s \in S$, the maps $\varphi_i(s) : C_i \rightarrow C_i$ define a natural transformation $F \circ G \rightarrow F \circ G$ mapped by the limit functor to $\lim_i (\varphi_i(s))$. It is easily seen that $(\lim_i C_i, \lim_i \varphi_i)$ where $(\lim_i \varphi_i)(s) = \lim_i (\varphi_i(s))$ is a limit of G . \square

When \mathcal{C} is abelian, the same holds for \mathcal{C}_S , and in this case $\mathcal{C}_S \rightarrow \mathcal{C}$ is exact. From now on we will only use the functor $(-)_S$ for abelian categories.

Assume now that $R \rightarrow S$ is a ring morphism between coherent commutative rings such that S is finitely presented as R -module. The functors $(S \otimes_R -) : \mathcal{C} \rightarrow \mathcal{C}$ and $\text{Hom}_R(S, -) : \mathcal{C} \rightarrow \mathcal{C}$ of Section 2.3 yield functors

$$(S \otimes_R -) : \mathcal{C} \rightarrow \mathcal{C}_S$$

and

$$\text{Hom}_R(S, -) : \mathcal{C} \rightarrow \mathcal{C}_S.$$

It is easily seen that

Proposition 4.3. *The functors $(S \otimes_R -)$ and $\text{Hom}_R(S, -)$ are respectively left and right adjoint to $\mathcal{C}_S \rightarrow \mathcal{C}$. In particular, $(S \otimes_R -)$ preserves projectives and $\text{Hom}_R(S, -)$ preserves injectives. \square*

The next result gives a connection between the functors $S \otimes_R (-)$ and $(-)_S$.

Proposition 4.4. *Let \mathfrak{a} be an essentially small R -linear category. Then*

(1) *there is a commutative diagram*

$$\begin{array}{ccc} \text{Mod}(S \otimes_R \mathfrak{a}) & \xrightarrow{\cong} & \text{Mod}(\mathfrak{a})_S \\ \alpha \downarrow & & \downarrow \beta \\ \text{Mod}(\mathfrak{a}) & \xlongequal{\quad} & \text{Mod}(\mathfrak{a}) \end{array}$$

where α is dual to $\mathfrak{a} \rightarrow S \otimes_R \mathfrak{a}$ and β is the forgetful functor;

(2) *the left and right adjoints to the forgetful functor $\text{Mod}(\mathfrak{a})_S \rightarrow \text{Mod}(\mathfrak{a})$ are computed pointwise, i.e.*

$$\begin{aligned} (S \otimes_R F)(A) &= S \otimes_R FA \\ \text{Hom}_R(S, F)(A) &= \text{Hom}_R(S, F(A)) \end{aligned}$$

In particular $S \otimes_R \mathfrak{a}(A, -) = (S \otimes_R \mathfrak{a})(A, -)$. \square

We have the following relation between $\text{Ind}(-)$ and $(-)_S$:

Proposition 4.5. *Assume that \mathcal{C} is essentially small. The obvious functor*

$$H : \text{Ind}(\mathcal{C}_S) \rightarrow (\text{Ind}(\mathcal{C}))_S$$

is an equivalence.

Proof. Consider the following commutative diagram with obvious maps.

$$\begin{array}{ccccc} \mathcal{C} & \longrightarrow & \text{Ind}(\mathcal{C}) & & \\ \uparrow F & & \uparrow \text{Ind}(F) & \swarrow & \\ \mathcal{C}_S & \longrightarrow & \text{Ind}(\mathcal{C}_S) & \xrightarrow{H} & (\text{Ind}(\mathcal{C}))_S \end{array}$$

Since F is faithful, the same holds for $\text{Ind}(F)$ and thus for H .

To prove that H is full, consider a map $(f_i)_i : \text{colim}_i(C_i, \varphi_i) \rightarrow \text{colim}_j(D_j, \psi_j)$ with $f_i : C_i \rightarrow D_{j_i}$ maps in \mathcal{C} such that $(\psi_j)_j \circ (f_i)_i = (f_i)_i \circ (\varphi_i)_i$ in $\text{Ind}(\mathcal{C})$. This means that for every i we obtain a diagram

$$\begin{array}{ccccc} C_i & \xrightarrow{f_i} & D_{j_i} & \xrightarrow{g_i} & D_{k_i} \\ \varphi_i \downarrow & & \downarrow \psi_{j_i} & & \downarrow \psi_{k_i} \\ C_i & \xrightarrow{f_i} & D_{j_i} & \xrightarrow{g_i} & D_{k_i} \end{array}$$

in which the right square commutes and composition with g_i makes the left square commute. It follows that the maps $g_i \circ f_i$ belong to \mathcal{C}_S and $(f_i)_i = (g_i \circ f_i)_i$.

To prove that H is essentially surjective, consider an object C of $(\text{Ind}(\mathcal{C}))_S$ (We omit the S -action φ in our notation). φ induces an $(\text{Ind}(\mathcal{C}))_S$ -epimorphism $S \otimes_R C \rightarrow C$ with kernel K , for which we obtain another $(\text{Ind}(\mathcal{C}))_S$ -epimorphism $S \otimes_R K \rightarrow K$. It follows that C is isomorphic to the cokernel of $S \otimes_R K \rightarrow S \otimes_R C$.

Writing $C = \operatorname{colim}_i C_i$ as a filtered colimit, we see that $S \otimes_R C = \operatorname{colim}_i (S \otimes_R C_i)$ belongs to $\operatorname{Ind}(\mathcal{C}_S)$ and the same holds for $S \otimes_R K$. Since $\operatorname{Ind}(F)$ is exact, the cokernel of the map belongs to $\operatorname{Ind}(\mathcal{C}_S)$, as we wanted. \square

Next we consider finitely presented objects.

Proposition 4.6. *Assume that in \mathcal{C} small filtered colimits are exact. If (C, φ) is finitely presented (resp finitely generated) in \mathcal{C}_S , then C is finitely presented (resp. finitely generated) in \mathcal{C} . The obvious functor*

$$H : \operatorname{Fp}(\mathcal{C}_S) \longrightarrow (\operatorname{Fp}(\mathcal{C}))_S$$

is an isomorphism.

Proof. Consider a finitely presented (C, φ) in \mathcal{C}_S and a filtered colimit $\operatorname{colim}_i C_i$ in \mathcal{C} . Making use of Corollary 2.8 and the fact that $\mathcal{C}_S \rightarrow \mathcal{C}$ reflects colimits, we may compute

$$\begin{aligned} \mathcal{C}(C, \operatorname{colim}_i C_i) &= \mathcal{C}_S((C, \varphi), \operatorname{Hom}_R(S, \operatorname{colim}_i C_i)) \\ &= \mathcal{C}_S((C, \varphi), \operatorname{colim}_i \operatorname{Hom}_R(S, C_i)) \\ &= \operatorname{colim}_i \mathcal{C}_S((C, \varphi), \operatorname{Hom}_R(S, C_i)) \\ &= \operatorname{colim}_i \mathcal{C}(C, C_i), \end{aligned}$$

hence C is finitely presented in \mathcal{C} . Now consider the following commutative diagram with obvious maps.

$$\begin{array}{ccccc} \mathcal{C} & \longrightarrow & \operatorname{Fp}(\mathcal{C}) & & \\ \uparrow F & & \uparrow \operatorname{Fp}(F) & \swarrow & \\ \mathcal{C}_S & \longrightarrow & \operatorname{Fp}(\mathcal{C}_S) & \xrightarrow{H} & (\operatorname{Fp}(\mathcal{C}))_S \end{array}$$

H is readily seen to be fully faithful and injective on objects, so it remains to show that it is surjective on objects. Consider (C, φ) in $(\operatorname{Fp}(\mathcal{C}))_S$ and a filtered colimit $\operatorname{colim}_i (C_i, \varphi_i) = (\operatorname{colim}_i C_i, \operatorname{colim}_i \varphi_i)$ in \mathcal{C}_S . The canonical map

$$\operatorname{colim}_i \mathcal{C}_S((C, \varphi), (C_i, \varphi_i)) \longrightarrow \mathcal{C}_S((C, \varphi), (\operatorname{colim}_i C_i, \operatorname{colim}_i \varphi_i))$$

is obviously injective. We finish the proof by showing that it is surjective. If $f : C \rightarrow \operatorname{colim}_i C_i$ defines a map in the codomain, f factors over some $f_i : C \rightarrow C_i$ in \mathcal{C} . For each $s \in S$, there is a diagram

$$\begin{array}{ccccc} C & \longrightarrow & C_i & \longrightarrow & \operatorname{colim}_i C_i \\ \varphi(s) \downarrow & & \downarrow \varphi_i(s) & & \downarrow \operatorname{colim}_i \varphi_i(s) \\ C & \longrightarrow & C_i & \longrightarrow & \operatorname{colim}_i C_i \end{array}$$

in which the right hand square is not necessarily commutative. Selecting finitely many generators s_k of S over R , and using the fact that C is finitely presented in \mathcal{C} and that the colimit is filtered, we can find $C_i \rightarrow C_j$ such that the right hand square with C_j instead of C_i commutes for every generator s_k and hence also for every $s \in S$. \square

We mention the familiar change of rings spectral sequences.

Proposition 4.7. *Let $X \in \text{mod}(S)$, $C \in \mathcal{C}$, $A \in \mathcal{C}_S$. Assume that \mathcal{C} is flat. There are first quadrant spectral sequences*

$$(4.1) \quad E_2^{pq} : \text{Ext}_{\mathcal{C}_S}^p(A, \text{Ext}_R^q(S, C)) \Rightarrow \text{Ext}_{\mathcal{C}}^n(A, C)$$

$$(4.2) \quad E_2^{pq} : \text{Ext}_S^p(X, \text{Ext}_R^q(S, C)) \Rightarrow \text{Ext}_R^n(X, C)$$

Proof. By enlarging our universe we may assume that \mathcal{C} is small. Then we replace \mathcal{C} by $\text{Ind}(\mathcal{C})$ in order to have a category with enough injectives. By Proposition 3.6 this enlarged \mathcal{C} is still flat. Furthermore by Proposition 3.5 $\text{Ext}_R^q(S, -)$ is the derived functor of $\text{Hom}_R(S, -)$.

We have

$$(4.3) \quad \text{Hom}_{\mathcal{C}_S}(A, \text{Hom}_R(S, C)) = \text{Hom}_{\mathcal{C}}(A, C)$$

$$(4.4) \quad \text{Hom}_S(X, \text{Hom}_R(S, C)) = \text{Hom}_R(X, C)$$

Now $\text{Hom}_R(S, -)$ preserves injectives and (4.4) shows that $\text{Hom}_R(S, -)$ also preserves coflat objects. So $\text{Hom}_R(S, -)$ sends injectives to acyclic objects for $\text{Hom}_{\mathcal{C}_S}(A, -)$ and $\text{Hom}_S(X, -)$.

Thus (4.1)(4.2) are just the Grothendieck spectral sequences [11] associated to (4.3) (4.4). \square

Here are some properties that are preserved under base change.

Proposition 4.8. (1) *If \mathcal{C} has enough injectives, the same holds for \mathcal{C}_S ;*
 (2) *if $(G_i)_i$ is a set of generators of \mathcal{C} , $(S \otimes_R G_i)_i$ is a set of generators of \mathcal{C}_S ;*
 (3) *if \mathcal{C} is Grothendieck, the same holds for \mathcal{C}_S ;*
 (4) *if \mathcal{C} is a locally coherent Grothendieck category, the same holds for \mathcal{C}_S ;*
 (5) *if \mathcal{C} is flat over R , then \mathcal{C}_S is flat over S .*

Proof. Only (5) is not entirely clear from the above discussion. Consider $C \in \mathcal{C}_S$ and $X \in \text{mod}(S)$. Take a \mathcal{C} -monomorphism $m : C \rightarrow C'$ with $\text{Ext}_R^1(X, m) = 0$. Then $\text{Hom}_R(S, m) : \text{Hom}_R(S, C) \rightarrow \text{Hom}_R(S, C')$ is a monomorphism. The spectral sequence (4.2) yields monomorphism $\text{Ext}_S^1(X, \text{Hom}_R(S, C)) \rightarrow \text{Ext}_S^1(X, C)$ natural in C and hence $\text{Ext}_S^1(X, \text{Hom}_R(S, m)) = 0$. The composition of monomorphisms $C \rightarrow \text{Hom}_R(S, C) \rightarrow \text{Hom}_R(S, C')$ is the required effacing of $\text{Ext}_S^1(X, C)$ in \mathcal{C}_S . \square

5. DEFORMATIONS OF ABELIAN CATEGORIES

In this section we introduce deformations of pre-additive and abelian categories. Basically these are lifts along the functors $S \otimes_R (-)$ and $(-)_S$ for a homomorphism of commutative rings $\theta : R \rightarrow S$. In the sequel we will assume that θ is surjective with nilpotent kernel but this is not necessary for the basic definitions.

We start with the pre-additive case.

Definition 5.1. For an S -linear category \mathfrak{b} , an R -deformation of \mathfrak{b} is an R -linear category \mathfrak{a} together with an R -linear functor $\mathfrak{a} \rightarrow \overline{\mathfrak{b}}$ that induces an equivalence $S \otimes_R \mathfrak{a} \rightarrow \mathfrak{b}$. A deformation $\mathfrak{a} \rightarrow \overline{\mathfrak{b}}$ will often be denoted by $\mathfrak{a} \rightarrow \mathfrak{b}$ or simply by \mathfrak{a} . If $S \otimes_R \mathfrak{a} \rightarrow \mathfrak{b}$ is an isomorphism, the deformation is called *strict*. If \mathfrak{a} is flat over R , the deformation is called *flat*.

Now we consider the abelian case.

Definition 5.2. For an *abelian* S -linear category \mathcal{D} , an R -deformation of \mathcal{D} is an *abelian* R -linear category \mathcal{C} together with an R -linear functor $\overline{\mathcal{D}} \rightarrow \mathcal{C}$ that induces an equivalence $\mathcal{D} \rightarrow \mathcal{C}_S$. A deformation $\overline{\mathcal{D}} \rightarrow \mathcal{C}$ will often be denoted by $\mathcal{D} \rightarrow \mathcal{C}$ or simply by \mathcal{C} . If $\mathcal{D} \rightarrow \mathcal{C}_S$ is an isomorphism, the deformation is called *strict*. If \mathcal{C} is flat over R (as an abelian category), the deformation is called *flat*.

Remark 5.3. In case of possible confusion we refer to the deformations of Definition 5.1 as *linear deformations* and to the deformations of Definition 5.2 as *abelian deformations*.

We will now assume that $\theta : R \rightarrow S$ is a homomorphism between coherent commutative rings such that S is a finitely presented R -module. Then the left and right adjoint to an abelian deformation $\mathcal{D} \rightarrow \mathcal{C}$ exist and we will continue to denote them by $S \otimes_R -$ and $\mathrm{Hom}_R(S, -)$.

Rephrasing the results of Sections 3,4 in terms of deformations we deduce that (flat) deformations are preserved under some natural constructions.

Proposition 5.4. *Let \mathfrak{b} be an essentially small S -linear category. An essentially small linear R -deformation $\mathfrak{a} \rightarrow \mathfrak{b}$ induces an abelian deformation $\mathrm{Mod}(\mathfrak{b}) \rightarrow \mathrm{Mod}(\mathfrak{a})$. Moreover flatness of these deformations is equivalent.*

Proof. We may assume that $\mathfrak{b} = S \otimes_R \mathfrak{a}$. Then the result follows from Propositions 3.7 and 4.4. \square

Proposition 5.5. *Every abelian R -deformation $\mathcal{D} \rightarrow \mathcal{C}$ between essentially small categories induces an abelian R -deformation $\mathrm{Ind}(\mathcal{D}) \rightarrow \mathrm{Ind}(\mathcal{C})$. Moreover flatness of these deformations is equivalent.*

Proof. We may assume $\mathcal{D} = \mathcal{C}_S$. Then the result follows from Propositions 3.6 and 4.5. \square

Proposition 5.6. *A deformation $\mathcal{D} \rightarrow \mathcal{C}$ of locally coherent Grothendieck categories induces a deformation $\mathrm{Fp}(\mathcal{D}) \rightarrow \mathrm{Fp}(\mathcal{C})$. Moreover flatness of these deformations is equivalent.*

Proof. We may assume $\mathcal{D} = \mathcal{C}_S$. Then the result follows from Propositions 3.6 and 4.6(4). \square

6. PRESERVATION OF PROPERTIES UNDER NILPOTENT DEFORMATIONS

In this chapter we investigate some categorical/homological properties that are preserved under flat, *nilpotent* (see below) abelian deformation. More precisely, we will show that the following properties of an abelian category \mathcal{D} lift to a flat nilpotent deformation:

- (1) \mathcal{D} is essentially small;
- (2) \mathcal{D} has enough injectives;
- (3) \mathcal{D} is a Grothendieck category;
- (4) \mathcal{D} is a locally coherent Grothendieck category.

lift to a flat nilpotent deformation.

6.1. Nilpotent deformations. As usual $\theta : R \rightarrow S$ is a morphism between coherent commutative rings such that S is finitely presented over R . In order to lift properties from an S -linear (abelian) category to an R -deformation, we obviously need some further assumptions on the ring map θ . First of all, we will assume that θ is surjective. In this case, for an abelian R -linear \mathcal{C} , the forgetful functor $\mathcal{C}_S \rightarrow \mathcal{C}$ is fully faithful. Put $I = \ker \theta$. The hypotheses imply that I is finitely presented.

Next, we will assume that the kernel of θ is nilpotent.

Definition 6.1. The R -deformations (linear or abelian) are called *nilpotent* provided the ideal I is nilpotent, i.e. $I^n = 0$ for some n . If $I^n = 0$ the deformation is called (*nilpotent*) of order n .

Remark 6.2. If \mathcal{C} a nilpotent deformation of \mathcal{D} (linear or abelian), then clearly there exist categories $\mathcal{D} = \mathcal{C}_0, \mathcal{C}_1, \dots, \mathcal{C}_k = \mathcal{C}$ for certain k such that \mathcal{C}_{i+1} is a nilpotent deformation of order 2 of \mathcal{C}_i . If \mathcal{C} is flat the we may assume that the $(\mathcal{C}_i)_i$ are flat as well.

Proposition 6.3. (*“Nakayama”*) Consider a nilpotent deformation $\mathcal{D} \rightarrow \mathcal{C}$ and $C \in \mathcal{C}$. If either $S \otimes_R C = 0$ or $\text{Hom}_R(S, C) = 0$, then $C = 0$.

Proof. If $S \otimes_R C = 0$ then $C = IC$ (§2.3). Since I is nilpotent this implies $C = 0$. The case $\text{Hom}_R(S, C) = 0$ is similar. \square

Proposition 6.4. Consider a nilpotent R -deformation $\mathcal{D} \rightarrow \mathcal{C}$. The functor $\text{Hom}_R(S, -) : \mathcal{C} \rightarrow \mathcal{D}$ reflects monomorphisms.

Proof. This readily follows from Proposition 6.3 using kernels. \square

6.2. Preservation of size. In this section we temporarily drop the assumption that our categories are automatically \mathcal{U} -categories and that the base rings R, S are \mathcal{U} -small.

We show that nilpotent deformations behave well with respect to size matters.

Lemma 6.5. For $M \in \text{Mod}(R)$,

$$|M| \leq \sup\{|S \otimes_R M|, |\mathbb{N}|\}.$$

Proof. This follows by considering the I -adic filtration on M . \square

For a category \mathcal{C} , the *skeleton* $\text{Sk}(\mathcal{C})$ of \mathcal{C} is the set of all isomorphism classes of \mathcal{C} -objects.

Lemma 6.6. Consider an S -linear category \mathfrak{b} and let $f : \mathfrak{a} \rightarrow \mathfrak{b}$ be a nilpotent R -deformation of \mathfrak{b} . Suppose α is an infinite cardinal such that $|\text{Sk}(\mathfrak{b})| \leq \alpha$ and for all $B, B' \in \text{Ob}(\mathfrak{b})$ one has $|\mathfrak{b}(B, B')| \leq \alpha$. Then the analogous property holds for \mathfrak{a} .

Proof. This follows from the following observations:

- (1) By Lemma 6.5, $|\mathfrak{a}(A, A')| \leq \sup\{|\mathfrak{b}(f(A), f(A'))|, |\mathbb{N}|\}$ for $A, A' \in \mathfrak{a}$;
- (2) By Proposition A.1, $|\text{Sk}(\mathfrak{a})| = |\text{Sk}(\mathfrak{b})|$. \square

Proposition 6.7. Suppose S is \mathcal{U} -small. Consider an S -linear category \mathfrak{b} and let $f : \mathfrak{a} \rightarrow \mathfrak{b}$ be a nilpotent R -deformation of \mathfrak{b} . If \mathfrak{b} is a \mathcal{U} -category (resp. essentially \mathcal{U} -small), the same holds for \mathfrak{a} .

Proof. Immediate from (the proof of) Lemma 6.6. \square

We want to prove a similar result for abelian deformations. We prove an analogue of Lemma 6.6. Note that the result we prove is more general than what we immediately need but its more general form will be used afterwards.

Lemma 6.8. *Suppose S is \mathcal{U} -small. Consider a flat deformation $\mathcal{D} \rightarrow \mathcal{C}$ of abelian categories which is of order two. Let \mathcal{D}' be a Serre subcategory of \mathcal{D} (i.e. a full subcategory closed under subquotients and extensions) and let \mathcal{C}' be the Serre subcategory of \mathcal{C} generated by \mathcal{D}' (i.e. the full subcategory of \mathcal{C} of objects C with $S \otimes_R C$ (hence also IC) $\in \mathcal{D}'$, see also Proposition 7.5).*

Suppose α is an infinite cardinal such that $|\mathrm{Sk}(\mathcal{D}')| \leq \alpha$ and for all $D, D' \in \mathrm{Ob}(\mathcal{D}')$ one has $|\mathcal{D}(D, D')| \leq \alpha$. Then the analogous property holds for \mathcal{C}' .

Proof. Choose C, C' . Filtering C, C' by the I -adic filtration we immediately deduce

$$(6.1) \quad |\mathcal{C}(C, C')| \leq |\mathcal{D}(S \otimes_R C, IC')| \cdot |\mathcal{D}(S \otimes_R C, S \otimes_R C')|$$

This implies $|\mathcal{C}(C, C')| \leq \alpha$

Concerning $|\mathrm{Sk}(\mathcal{C}')|$, we make the following observations:

- (1) An object in \mathcal{C}' is up to isomorphism determined by a triple (A, B, e) where $A, B \in \mathrm{Ob}(\mathcal{D}')$ and $e \in \mathrm{Ext}_{\mathcal{C}}^1(A, B)$.
- (2) The spectral sequence (4.1) yields the bound:

$$|\mathrm{Ext}_{\mathcal{C}}^1(A, B)| \leq |\mathrm{Ext}_{\mathcal{D}}^1(A, B)| |\mathrm{Hom}_{\mathcal{D}}(A, \mathrm{Ext}_R^1(S, B))|$$

- (3) An element of $\mathrm{Ext}_{\mathcal{D}}^1(A, B)$ may be represented by a triple (C, u, v) where $C \in \mathrm{Ob}(\mathcal{D}')$ and $u \in \mathcal{D}(A, C)$, $v \in \mathcal{D}(C, B)$ are such that (u, v) constitutes a short exact sequence.

The bound $|\mathrm{Sk}(\mathcal{C}')| \leq \alpha$ now easily follows. \square

Proposition 6.9. *Suppose S is \mathcal{U} -small. Consider a flat nilpotent deformation $\mathcal{D} \rightarrow \mathcal{C}$ of abelian categories. If \mathcal{D} is a \mathcal{U} -category (resp. an essentially \mathcal{U} -small category), then so is \mathcal{C} .*

Proof. We may assume that the deformation is of order two. The case that \mathcal{D} is essentially \mathcal{U} -small follows directly from Lemma 6.8. For the case that \mathcal{D} is a \mathcal{U} -category we invoke (6.1) to obtain that every $|\mathcal{C}(C, C')|$ is bounded by the cardinality of an element of \mathcal{U} . \square

6.3. Lifting of objects. An important tool in the study of deformations is the lifting of objects along the functors $(S \otimes_R -)$ and $\mathrm{Hom}_R(S, -)$. In this section we state some results on lifting that will be used afterwards. We start with the following definition.

Definition 6.10. Consider a functor $H : \mathcal{C} \rightarrow \mathcal{D}$ and an object D of \mathcal{D} . A *lift* of D along H is an object C of \mathcal{C} and an isomorphism $D \cong H(C)$. Lifts of D , together with the obvious morphisms, constitute a groupoid $L(H, D)$.

Remark 6.11. If $H : \mathcal{C} \rightarrow \mathcal{D}$ is right adjoint to a functor $F : \mathcal{D} \rightarrow \mathcal{C}$, a lift of D along H can be represented by a map $F(D) \rightarrow C$. If H is left adjoint to $G : \mathcal{D} \rightarrow \mathcal{C}$, a lift of D along H can be represented by a map $C \rightarrow G(D)$.

Consider a flat deformation $\mathcal{D} \rightarrow \mathcal{C}$. The restriction $\mathrm{Hom}_R^{cf}(S, -)$ of $\mathrm{Hom}_R(S, -)$ to coflat objects yields a groupoid

$$L^{cf}(D) = L(\mathrm{Hom}_R^{cf}(S, -), D)$$

for every coflat D in \mathcal{D} . Analogously, the restriction $(S \otimes_R^f -)$ of $(S \otimes_R -)$ to flat objects yields a groupoid

$$L^f(D) = L((S \otimes_R^f -), D)$$

for every flat D in \mathcal{D} .

We state the following theorem without proof. The theorem is a special case of an obstruction theory for derived lifting, which will be published separately [16].

Theorem 6.12. *Consider a flat nilpotent R -deformation $\mathcal{D} \rightarrow \mathcal{C}$ of order 2 and a coflat object D of \mathcal{D} . There is an obstruction*

$$o(D) \in \text{Ext}_{\mathcal{D}}^2(\text{Hom}_S(I, D), D)$$

satisfying

$$L^{cf}(D) \neq \emptyset \iff o(D) = 0.$$

over

$$\text{Ext}_{\mathcal{D}}^1(\text{Hom}_S(I, D), D).$$

The following proposition shows that in a certain sense, the conditions of Theorem 6.12 can itself be lifted under deformation.

Proposition 6.13. *Consider a flat nilpotent deformation $\mathcal{D} \rightarrow \mathcal{C}$ and coflat \mathcal{C} -objects \overline{D} and \overline{E} with $\text{Hom}_R(S, \overline{D}) = D$ and $\text{Hom}_R(S, \overline{E}) = E$. Suppose that for certain $i \geq 0$ we have that $\text{Ext}_{\mathcal{D}}^i(\text{Hom}_S(X, D), E) = 0$ for all $X \in \text{mod}(S)$. Then $\text{Ext}_{\mathcal{C}}^i(\text{Hom}_R(X, \overline{D}), \overline{E}) = 0$ for all $X \in \text{mod}(R)$.*

Proof. We may consider $\mathcal{C}_S \rightarrow \mathcal{C}$ of order 2. By Proposition 4.7,

$$\text{Ext}_{\mathcal{C}}^i(\text{Hom}_R(Y, \overline{D}), \overline{E}) = \text{Ext}_{\mathcal{D}}^i(\text{Hom}_S(Y, D), E) = 0$$

for all $Y \in \text{mod}(S)$. For $X \in \text{mod}(R)$, it suffices to consider $0 \rightarrow IX \rightarrow X \rightarrow S \otimes_R X \rightarrow 0$ and the corresponding $0 \rightarrow \text{Hom}_R(S \otimes_R X, \overline{D}) \rightarrow \text{Hom}_R(X, \overline{D}) \rightarrow \text{Hom}_R(IX, \overline{D}) \rightarrow 0$. The result for X then follows from the result for $Y = IX$ and $Y = S \otimes_R X$ via the long exact Ext-sequence. \square

Corollary 6.14. *Consider a flat nilpotent deformation $\mathcal{D} \rightarrow \mathcal{C}$ and a coflat \mathcal{D} -object D . If $\text{Ext}_{\mathcal{D}}^2(\text{Hom}_S(X, D), D) = 0$ for all $X \in \text{mod}(S)$, then $\text{Sk}(L^{cf}(D)) \neq \emptyset$. If $\text{Ext}_{\mathcal{D}}^2(\text{Hom}_S(X, D), D) = \text{Ext}_{\mathcal{D}}^1(\text{Hom}_S(X, D), D) = 0$, then $\text{Sk}(L^{cf}(D))$ contains precisely one element.*

Proof. This follows from Propositions 4.8, 6.13 and Theorem 6.12. \square

Corollary 6.15. *Consider a flat nilpotent deformation $\mathcal{D} \rightarrow \mathcal{C}$ and a \mathcal{D} -injective D . Then $\text{Sk}(L^{cf}(D))$ contains precisely one lift $D \rightarrow \mathcal{C}$ and \mathcal{C} is a \mathcal{C} -injective.*

Proof. We may consider $\mathcal{C}_S \rightarrow \mathcal{C}$ of order 2. The first part of the statement follows from Theorem 6.12. To prove that \mathcal{C} is injective, first notice that by Proposition 4.7, $\text{Ext}_{\mathcal{C}}^i(A, C) = 0$ for all $A \in \mathcal{C}_S$. Finally, for $A \in \mathcal{C}$, it suffices to consider the long exact Ext sequence associated to $0 \rightarrow IA \rightarrow A \rightarrow S \otimes_R A \rightarrow 0$. \square

6.4. Deformations of categories with enough injectives. The results of the previous section allow us to prove the following theorem:

Theorem 6.16. *Consider a flat nilpotent deformation $\mathcal{D} \rightarrow \mathcal{C}$. If \mathcal{D} has enough injectives then so does \mathcal{C} .*

Proof. We may consider $\mathcal{C}_S \rightarrow \mathcal{C}$ and an object C in \mathcal{C} . Take a \mathcal{C}_S -monomorphism $m : \text{Hom}_R(S, C) \rightarrow I$ to a \mathcal{C}_S -injective I such that there is a lift \bar{I} with $\text{Hom}_R(S, \bar{I}) = I$. By Corollary 6.15, \bar{I} is injective and so m can be lifted to a map $\bar{m} : C \rightarrow \bar{I}$ with $\text{Hom}_R(S, \bar{m}) = m$. By Proposition 6.4, \bar{m} is a \mathcal{C} -monomorphism. \square

This result is false for non-flat nilpotent deformations, as the following example shows.

Example 6.17. Let k be a field and let $S = k$, $R = k[\epsilon]$, $\epsilon^2 = 0$. Let $\mathcal{D} = \text{Mod}(k)$. Let κ be an infinite small cardinal and consider the following category:

$$\mathcal{C} = \{M \in \text{Mod}(R) \mid \dim_k IM \leq \kappa\}$$

It is easy to see that \mathcal{C} is full abelian subcategory of $\text{Mod}(R)$. Clearly $\mathcal{D} \rightarrow \mathcal{C}$ is a nilpotent deformation of order two.

A skeletal subcategory of $\text{Mod}(R)$ is given by the objects

$$V(\alpha, \beta) = k^{\oplus \alpha} \bigoplus k[\epsilon]^{\oplus \beta}$$

where α, β are small cardinals [23]. \mathcal{C} consists of those objects $V(\alpha, \beta)$ for which $\beta \leq \kappa$.

The objects $V(\alpha, \beta)$ with $\alpha \neq 0$ are not injective in \mathcal{C} since we may always replace one copy of k by $k[\epsilon]$ which yields a non-split extension.

Consider an object of the form $V(\alpha, \kappa)$ in \mathcal{C} with $\alpha > \kappa$ and assume that there is an injective object E in \mathcal{C} containing $V(\alpha, \kappa)$. Then clearly E must be of the form $V(\alpha', \kappa)$ with $\alpha' + \kappa \geq \alpha + \kappa$. In particular $\alpha' > \kappa$. But then E cannot be injective by the discussion in the previous paragraph. This contradiction shows that \mathcal{C} does not have enough injectives.

6.5. Deformations of Grothendieck categories. The aim of this section is to prove that the property of being a Grothendieck category is preserved under flat nilpotent deformation.

For a small pre-additive category \mathfrak{g} we write $\text{Pr}(\mathfrak{g}) = \text{Mod}(\mathfrak{g}^{\text{op}})$ for the category of (additive) presheaves on \mathfrak{g} . We recall the following version of the Gabriel-Popescu theorem.

Theorem 6.18. [21, 22, 15] *Let \mathcal{D} be a Grothendieck category and let \mathfrak{g} be a small generating subcategory of \mathcal{D} . There is a localization (i.e. a fully faithful functor with an exact left adjoint)*

$$\mathcal{D} \rightarrow \text{Pr}(\mathfrak{g}) : D \mapsto \mathcal{D}(-, D)$$

Conversely if \mathfrak{g} is a small pre-additive category and $\mathcal{D} \rightarrow \text{Pr}(\mathfrak{g})$ a localization, then \mathcal{D} is a Grothendieck category. Equivalence classes of such localizations are in one-one correspondence with so-called Gabriel (or additive) topologies on \mathfrak{g} .

Let \mathcal{D} be a Grothendieck category with a distinguished generator G . We define a cardinality $|\cdot|$ on $\text{Ob}(\mathcal{D})$ by

$$|C| = |\mathcal{D}(G, C)|,$$

where $|\mathcal{D}(G, C)|$ denotes the set theoretic cardinality of the set $\mathcal{D}(G, C)$. For a cardinal κ , \mathcal{D}_κ is the full subcategory of \mathcal{D} containing all \mathcal{D} -objects D with $|D| \leq \kappa$.

Example 6.19. If we take \mathcal{D} to be a module category $\mathbf{Mod}(A)$ over a ring A and we take $G = A$, the newly defined $|M|$ coincides with the set theoretic cardinality of M .

The following proposition lists the essential properties of the categories \mathcal{D}_κ .

Proposition 6.20. *If κ is a small cardinal, then \mathcal{D}_κ is essentially small. If κ is an infinite cardinal of the form 2^β , with $\beta \geq |G|$ then \mathcal{D}_κ is a Serre subcategory of \mathcal{D} .*

Proof. We prove first that \mathcal{D}_κ is essentially small. By Theorem 6.18 \mathcal{D}_κ is equivalent to a full subcategory of $\mathbf{Mod}(\mathcal{D}(G, G)^{\text{op}})_\kappa$. The latter obviously has small skeleton since the set of right $\mathcal{D}(G, G)$ -module structures on every $\kappa' \leq \kappa$ is small.

The fact that \mathcal{D}_κ is a Serre subcategory of \mathcal{D} is a consequence of Lemmas 6.21, 6.22 below which show that the ordinary properties of cardinalities of modules more or less hold for objects in Grothendieck categories. The only not entirely obvious fact is that \mathcal{D}_κ is closed under quotients under the stated hypotheses on κ . This follows from Lemma 6.22 and the following computation.

$$(2\kappa)^{|G|} = 2^{\beta \cdot |G|} = 2^\beta = \kappa \quad \square$$

Lemma 6.21. *For an exact sequence $0 \rightarrow A \rightarrow B \rightarrow C$ in \mathcal{D} ,*

- (1) $|A| \leq |B|$;
- (2) $|B| \leq |A| \cdot |C|$.

Proof. This follows from the exactness of $0 \rightarrow \mathcal{D}(G, A) \rightarrow \mathcal{D}(G, B) \rightarrow \mathcal{D}(G, C)$. \square

Lemma 6.22. *For an epimorphism $f : A \rightarrow B$,*

$$|B| \leq (2|A|)^{|G|}.$$

Proof. For a map $g : G \rightarrow B$, consider the pullback P_g of f and g and then an epimorphism $\bigoplus_{i \in I} G \rightarrow P_g$. The composition $(f_i)_{i \in I} : \bigoplus_{i \in I} G \rightarrow P_g \rightarrow G$ is an epimorphism. Put $J = \{f_i \mid i \in I\} \subset \mathcal{D}(G, G)$.

In this way we still have an epimorphism $(j)_{j \in J} : \bigoplus_{j \in J} G \rightarrow G$ which fits in a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ (a_j)_j \uparrow & & \uparrow g \\ \bigoplus_{j \in J} G & \xrightarrow{(j)_j} & G \end{array}$$

Conversely it is clear that g is uniquely determined by the data $J \subset \mathcal{D}(G, G)$ and $(a_j)_{j \in J} \in \mathcal{D}(G, A)^J$ which yields the required bound. \square

In the remainder of this section, we consider a flat nilpotent deformation $\mathcal{D} = \mathcal{C}_S \rightarrow \mathcal{C}$ of order 2 in which \mathcal{D} is a Grothendieck category with a fixed generator G defining a cardinality $|\cdot|$ on \mathcal{D} . We put $\kappa = 2^\beta$ for an infinite cardinal $\beta \geq |G|$. By Proposition 6.20, \mathcal{D}_κ is an essentially small Serre subcategory of \mathcal{D} . We let \mathcal{C}_κ be the Serre subcategory of \mathcal{C} generated by \mathcal{D}_κ . By Lemma 6.8 \mathcal{C}_κ is essentially small as well. Our aim is to show that \mathcal{C}_κ generates \mathcal{C} .

The following lemma gives us a general procedure for constructing generators for \mathcal{C} . It is more general than what we need but it will be reapplied in a slightly different setting afterwards (see Theorem 6.36).

Lemma 6.23. *Let $(G_i)_{i \in I}$ be a (not necessarily small) collection of generators of \mathcal{D} . Consider a collection of objects $(G_f)_f$, indexed over all non-zero \mathcal{C} -maps f with codomain in \mathcal{D} , obtained in the following manner:*

(6.2)

$$\begin{array}{ccccccccc}
 G_f & \xrightarrow{k} & G''_f & \xrightarrow{h'} & G' & \xrightarrow{1} & G' & \xrightarrow{g'} & A & \xrightarrow{f} & B \\
 & \searrow \gamma'' & \downarrow \gamma' & & \downarrow \gamma & & \downarrow \beta' & & \downarrow \beta & \nearrow f' & \\
 & & G_j & \xrightarrow{h} & S \otimes_R G' & \xrightarrow{\beta''} & G_i & \xrightarrow{g} & S \otimes_R A & &
 \end{array}$$

(P1) is between $G''_f \rightarrow G'$ and $G_j \rightarrow S \otimes_R G'$. (P2) is between $G' \rightarrow G_i$ and $S \otimes_R G' \rightarrow S \otimes_R A$.

g is chosen to make $f' \circ g \neq 0$, G' is the pullback of β and g , h is chosen to make $f' \circ g \circ \beta'' \circ h \neq 0$, G''_f is the pullback of γ and h , and k is any map such that $\gamma' = \gamma \circ k$ is an epimorphism. Then $(G_f)_f$ is a (not necessarily small) collection of generators for \mathcal{C} .

Proof. The G_f clearly satisfy the generator property with respect to the maps $f : A \rightarrow B$ with codomain in \mathcal{D} . We claim this is sufficient. Let $g : A \rightarrow C$ be a general map in \mathcal{C} . If the composition $A \xrightarrow{g} C \rightarrow S \otimes_R C$ is not-zero then we are done. If the composition is zero then we have a factorization $A \rightarrow IC \rightarrow C$ and we are done as well. \square

Lemma 6.24. *Suppose $I = \{*\}$, $G_* = G$ and let $(G_f)_f$ be constructed as in Lemma 6.23 with every k taken to be $1_{G''_f}$. Then for all f we have $G_f \in \mathcal{C}_\kappa$. In particular, \mathcal{C} has a small set of generators.*

Proof. Recall that the squares marked with (P*) in (6.2) are pullbacks. From (P1) it follows that we have to show that $IG' \in \mathcal{D}_\kappa$. From (P2) we obtain an exact sequence

$$0 \rightarrow IA \rightarrow G' \rightarrow G \rightarrow 0$$

Tensoring this sequence with S we obtain a commutative diagram with exact rows and columns

$$\begin{array}{ccccccc}
 & & & & 0 & & \\
 & & & & \uparrow & & \\
 \text{Tor}_1^R(S, G) & \longrightarrow & IA & \longrightarrow & S \otimes_R G' & \longrightarrow & G \longrightarrow 0 \\
 & & \parallel & & \uparrow & & \parallel \\
 0 & \longrightarrow & IA & \longrightarrow & G' & \longrightarrow & G \longrightarrow 0 \\
 & & & & \uparrow & & \\
 & & & & IG' & & \\
 & & & & \uparrow & & \\
 & & & & 0 & &
 \end{array}$$

Diagram chasing yields an epimorphism $\mathrm{Tor}_1^R(S, G) \longrightarrow IG'$ which finishes the proof. \square

Proposition 6.25. *Let \mathcal{E} be an abelian category with enough injectives. Put $\mathfrak{i} = \mathrm{Inj}(\mathcal{E})$. Consider the functor*

$$\Psi : \mathcal{E}^{\mathrm{op}} \longrightarrow \mathrm{Mod}(\mathfrak{i}) : C \longmapsto \mathcal{E}(C, -).$$

- (1) Ψ is exact, limit preserving and fully faithful;
- (2) Ψ induces an equivalence

$$\Psi' : \mathcal{E}^{\mathrm{op}} \longrightarrow \mathrm{mod}(\mathfrak{i})$$

Proof. Ψ is exact and limit preserving since all the functors $\mathcal{E}(-, I) : \mathcal{E}^{\mathrm{op}} \longrightarrow \mathbf{Ab}$ are exact and limit preserving.

To show that Ψ is fully faithful take $C, D \in \mathrm{Ob}(\mathcal{E})$. Choose an injective copresentation

$$0 \longrightarrow D \longrightarrow I_0 \longrightarrow I_1$$

Then we obtain a projective presentation of $\mathcal{E}(D, -)$ in $\mathrm{Mod}(\mathfrak{i})$

$$\mathfrak{i}(I_1, -) \longrightarrow \mathfrak{i}(I_0, -) \longrightarrow \mathcal{E}(D, -) \longrightarrow 0$$

and thus

$$\begin{aligned} \mathrm{Mod}(\mathfrak{i})(\mathcal{E}(D, -), \mathcal{E}(C, -)) &= \ker(\mathrm{Mod}(\mathfrak{i})(\mathfrak{i}(I_0, -), \mathcal{E}(C, -)) \longrightarrow \mathrm{Mod}(\mathfrak{i})(\mathfrak{i}(I_1, -), \mathcal{E}(C, -))) \\ &= \ker(\mathcal{E}(C, I_0) \longrightarrow \mathcal{E}(C, I_1)) \\ &= \mathcal{E}(C, D) \end{aligned}$$

Now consider $F \in \mathrm{mod}(\mathfrak{i})$. Then F has a presentation

$$\bigoplus_{i=1}^n \mathfrak{i}(I_i, -) \longrightarrow \bigoplus_{j=1}^m \mathfrak{i}(I_j, -) \longrightarrow F \longrightarrow 0,$$

hence it follows that F is isomorphic to $\mathcal{E}(C, -)$ with C defined by $0 \longrightarrow C \longrightarrow \bigoplus_{j=1}^m I_j \longrightarrow \bigoplus_{i=1}^n I_i$. This proves that Ψ' is essentially surjective. \square

Lemma 6.26. *\mathcal{C} has arbitrary small coproducts and they are exact.*

Proof. Besides our fixed universe \mathcal{U} we introduce a larger universe \mathcal{V} such that \mathcal{C} is \mathcal{V} -small.

By Theorem 6.16, \mathcal{C} has enough injectives and putting $\mathfrak{i} = \mathrm{Inj}(\mathcal{C})$, by Proposition 6.25 we have an exact, limit preserving functor $\Psi : \mathcal{C}^{\mathrm{op}} \longrightarrow \mathcal{V}\text{-Mod}(\mathfrak{i}) : C \longmapsto \mathcal{C}(C, -)$ inducing an equivalence $\Psi' : \mathcal{C}^{\mathrm{op}} \longrightarrow \mathcal{V}\text{-mod}(\mathfrak{i})$. We are to prove that $\mathcal{C}^{\mathrm{op}}$ has exact \mathcal{U} -small products. Since $\mathcal{V}\text{-Mod}(\mathfrak{i})$ has even exact \mathcal{V} -small products, it suffices to show that for a \mathcal{U} -small set of \mathcal{C} -objects $(C_i)_i$ the product $\prod_i \mathcal{C}(C_i, -)$ in $\mathcal{V}\text{-Mod}(\mathfrak{i})$ is finitely presented. Consider the short exact sequences in \mathcal{C}

$$0 \longrightarrow IC_i \longrightarrow C_i \longrightarrow S \otimes_R C_i \longrightarrow 0.$$

In $\mathcal{V}\text{-Mod}(\mathfrak{i})$ we can take the product of the exact sequences

$$0 \longrightarrow \mathcal{C}(S \otimes_R C_i, -) \longrightarrow \mathcal{C}(C_i, -) \longrightarrow \mathcal{C}(IC_i, -) \longrightarrow 0$$

to obtain an exact sequence

$$0 \longrightarrow \mathcal{C}\left(\prod_i S \otimes_R C_i, -\right) \longrightarrow \prod_i \mathcal{C}(C_i, -) \longrightarrow \mathcal{C}\left(\prod_i IC_i, -\right) \longrightarrow 0,$$

where we used that \mathcal{C}_S has small \mathcal{U} -coproducts and $\mathcal{C}_S \rightarrow \mathcal{C}$ preserves them. Then $\prod_i \mathcal{C}(C_i, -)$ is finitely presented in $\mathcal{V}\text{-Mod}(\mathfrak{i})$ as an extension of finitely presented objects. \square

The following proposition is the dual of [20, A.3.2].

Proposition 6.27. *Consider an abelian category \mathcal{E} with exact coproducts and any small category \mathcal{I} . The colimit functor*

$$\text{colim} : \text{Fun}(\mathcal{I}, \mathcal{E}) \longrightarrow \mathcal{E} : F \longrightarrow \text{colim}(F)$$

is trivially right exact and its left derived functors $L_i \text{colim}$ exist. Moreover if \mathcal{E}' is another abelian category with exact coproducts and $\phi : \mathcal{E} \rightarrow \mathcal{E}'$ is an exact functor preserving coproducts, the following diagram commutes:

$$\begin{array}{ccc} \text{Fun}(\mathcal{I}, \mathcal{E}) & \xrightarrow{(\phi \circ \cdot)} & \text{Fun}(\mathcal{I}, \mathcal{E}') \\ L_i \text{colim}_{\mathcal{E}} \downarrow & & \downarrow L_i \text{colim}_{\mathcal{E}'} \\ \mathcal{E} & \xrightarrow{\phi} & \mathcal{E}'. \end{array}$$

Lemma 6.28. *\mathcal{C} has exact filtered colimits.*

Proof. By Lemma 6.26 we know that filtered colimits exist in \mathcal{C} , so we are to prove that for a filtered category \mathcal{I} , the functor

$$\text{colim} : \text{Fun}(\mathcal{I}, \mathcal{C}) \longrightarrow \mathcal{C} : F \longrightarrow \text{colim}(F)$$

is exact. By Lemma 6.26 and Proposition 6.27, it suffices to prove that $L_1 \text{colim} = 0$. Consider $F : \mathcal{I} \rightarrow \mathcal{C}$. Using the exact sequences $0 \rightarrow IF(i) \rightarrow F(i) \rightarrow S \otimes_R F(i) \rightarrow 0$ and $\phi : \mathcal{C}_S \rightarrow \mathcal{C}$, we obtain an exact sequence

$$0 \rightarrow \phi \circ F' \rightarrow F \rightarrow \phi \circ F'' \rightarrow 0$$

for functors F' and $F'' : \mathcal{I} \rightarrow \mathcal{C}_S$. We obtain an exact sequence

$$L_1 \text{colim}(\phi \circ F') \rightarrow L_1 \text{colim}(F) \rightarrow L_1 \text{colim}(\phi \circ F''),$$

but since filtered colimits are exact in \mathcal{C}_S , it follows from Proposition 6.27 that both ends of the sequence and hence also the middle term $L_1 \text{colim}(F)$ are zero, which proves our assertion. \square

Theorem 6.29. *Consider a flat nilpotent deformation $\mathcal{D} \rightarrow \mathcal{C}$.*

If \mathcal{D} is a Grothendieck category, the same holds for \mathcal{C} .

Proof. It suffices to consider a flat deformation $\mathcal{C}_S \rightarrow \mathcal{C}$ of order 2. The result then immediately follows from Lemmas 6.24, 6.26 and 6.28. \square

6.6. Deformations of locally coherent Grothendieck categories. In this section we show that local coherence of a Grothendieck category is also preserved under flat nilpotent deformation. We begin with some preliminary results.

Definition 6.30. [13] An object C in an abelian category \mathcal{C} is called *fp-injective* provided that a monomorphism $X \rightarrow Y$ between finitely presented objects induces an epimorphism $\mathcal{C}(Y, C) \rightarrow \mathcal{C}(X, C)$.

Proposition 6.31. *A filtered colimit of fp-injectives is fp-injective.*

Proof. Let $\text{colim}_i C_i$ be a filtered colimit of fp-injectives and $0 \longrightarrow X \longrightarrow Y \longrightarrow Z \longrightarrow 0$ an exact sequence of finitely presented objects. Then all sequences $0 \longrightarrow \mathcal{C}(Z, C_i) \longrightarrow \mathcal{C}(Y, C_i) \longrightarrow \mathcal{C}(X, C_i) \longrightarrow 0$ are exact hence so is their filtered colimit $0 \longrightarrow \mathcal{C}(Z, \text{colim}_i C_i) \longrightarrow \mathcal{C}(Y, \text{colim}_i C_i) \longrightarrow \mathcal{C}(X, \text{colim}_i C_i) \longrightarrow 0$. \square

Proposition 6.32. [13, A.1] *In a locally coherent Grothendieck category \mathcal{C} , the following are equivalent for $C \in \mathcal{C}$:*

- (1) C is fp-injective;
- (2) $\text{Ext}^1(Z, C) = 0$ for every Z in $\text{Fp}(\mathcal{C})$;
- (3) $\text{Ext}^i(Z, C) = 0$ for all $i > 0$ for every Z in $\text{Fp}(\mathcal{C})$. \square

Lemma 6.33. *Let C be finitely presented in an abelian category \mathcal{C} with enough injectives. If filtered colimits of injectives are $\mathcal{C}(C, -)$ -acyclic, then $\text{Ext}_{\mathcal{C}}^n(C, -)$ preserves filtered colimits for $n \geq 0$.*

Proof. Consider a filtered colimit $\text{colim}_i A_i$. By [11], we can make a functorial choice of injective resolutions I_i of the objects A_i . Then $\text{colim}_i I_i$ is a $\mathcal{C}(C, -)$ -acyclic resolution of $\text{colim}_i A_i$, hence we may compute

$$\begin{aligned} \text{Ext}_{\mathcal{C}}^n(C, \text{colim}_i A_i) &= H^n(\mathcal{C}(C, \text{colim}_i I_i)) \\ &= H^n(\text{colim}_i \mathcal{C}(C, I_i)) \\ &= \text{colim}_i \text{Ext}_{\mathcal{C}}^n(C, A_i). \quad \square \end{aligned}$$

Proposition 6.34. *Let C be finitely presented in a locally coherent Grothendieck category \mathcal{C} . Then $\text{Ext}_{\mathcal{C}}^n(C, -)$ preserves filtered colimits for $n \geq 0$.*

Proof. This follows from Proposition 6.32 and Lemma 6.33. \square

Lemma 6.35. *Consider a flat nilpotent deformation of Grothendieck categories $\mathcal{C}_S \longrightarrow \mathcal{C}$ with \mathcal{C}_S locally coherent. For a finitely presented object C of \mathcal{C}_S , $\text{Ext}_{\mathcal{C}}^n(C, -)$ preserves filtered colimits for $n \geq 0$.*

Proof. By Lemma 6.33 it suffices to show that for a filtered colimit $\text{colim}_i I_i$ of injectives $\text{Ext}_{\mathcal{C}}^n(C, \text{colim}_i I_i) = 0$. Now by Corollary 2.8, $\text{colim}_i I_i$ is coflat. Consequently, we may use Propositions 4.7 and 6.34 to compute that

$$\begin{aligned} \text{Ext}_{\mathcal{C}}^n(C, \text{colim}_i I_i) &= \text{Ext}_{\mathcal{C}_S}^n(C, \text{Hom}_R(S, \text{colim}_i I_i)) \\ &= \text{Ext}_{\mathcal{C}_S}^n(C, \text{colim}_i \text{Hom}_R(S, I_i)) \\ &= \text{colim}_i \text{Ext}_{\mathcal{C}_S}^n(C, \text{Hom}_R(S, I_i)) \\ &= \text{colim}_i \text{Ext}_{\mathcal{C}}^n(C, I_i) = 0. \quad \square \end{aligned}$$

We are now able to prove the main theorem.

Theorem 6.36. *Consider a flat nilpotent deformation $\mathcal{D} \longrightarrow \mathcal{C}$ of Grothendieck categories. If \mathcal{D} is locally coherent, the same holds for \mathcal{C} .*

Proof. We may consider $\mathcal{C}_S \longrightarrow \mathcal{C}$ of order 2. Let $(G_i)_i$ be a small set of coherent generators of \mathcal{C}_S . We want to carry out the construction of Lemma 6.23 for specific maps k . Suppose we have constructed the objects G_f'' . Since \mathcal{C}_S is locally coherent, IG' is a small filtered colimit $IG' = \text{colim}_j F_j$ of coherent objects F_j of \mathcal{C}_S . Every

canonical map $F_i \rightarrow IG'$ induces a map $\text{Ext}_{\mathcal{C}}^1(G_j, F_i) \rightarrow \text{Ext}_{\mathcal{C}}^1(G_j, IG')$ mapping an extension $0 \rightarrow F_i \rightarrow E \rightarrow G_j \rightarrow 0$ to the pullback

$$\begin{array}{ccc} F_i & \longrightarrow & IG' \\ \downarrow & & \downarrow \\ E & \longrightarrow & P \\ & \searrow & \downarrow \\ & & G_j \end{array}$$

By Lemma 6.35, $\text{Ext}_{\mathcal{C}}^1(G_j, -)$ preserves filtered colimits, hence

$$\text{Ext}_{\mathcal{C}}^1(G_j, IG') = \text{colim}_i \text{Ext}_{\mathcal{C}}^1(G_j, F_i).$$

It then follows from the construction of this filtered colimit that the extension

$$0 \longrightarrow IG' \xrightarrow{\delta'} G_f'' \xrightarrow{\gamma'} G_j \longrightarrow 0$$

is in the image of one of the maps $\text{Ext}_{\mathcal{C}}^1(G_j, F_i) \rightarrow \text{Ext}_{\mathcal{C}}^1(G_j, IG')$. Hence we obtain an extension

$$0 \longrightarrow F_i \xrightarrow{\delta''} G_f \xrightarrow{\gamma''} G_j \longrightarrow 0$$

and a map $k : G_f \rightarrow G_f''$ as in Lemma 6.23. Hence $(G_f)_f$ is a set of generators for \mathcal{C} . Now (see §2.2) $\mathbf{Fp}(\mathcal{C}_S)$ has a small skeleton and since \mathcal{C} has small Yoneda Ext groups, it follows that the image of $(G_f)_f$ is small. Using Proposition 4.6, it is readily seen that the objects G_i and F_i are coherent in \mathcal{C} . It then follows that all objects of $(G_f)_f$ are finitely generated, hence \mathcal{C} is locally finitely generated. But then all objects of $(G_f)_f$ are finitely presented, hence \mathcal{C} is locally finitely presented. Finally all objects of $(G_f)_f$ are coherent hence it follows that \mathcal{C} is locally coherent, as we set out to prove. \square

7. DEFORMATION AND LOCALIZATION

7.1. Statement of the main results. A fully faithful functor $i : \mathcal{L} \rightarrow \mathcal{C}$ between Grothendieck categories is called a *localization* if it has an exact left adjoint. If in addition i is an embedding of a full subcategory closed under isomorphisms then we call i a strict localization. In this section, we study the compatibility of localization with deformations.

Below we prove the following result.

Theorem 7.1. *Let $\mathcal{C}_S \rightarrow \mathcal{C}$ be a nilpotent deformation of Grothendieck categories. Then*

$$\mathcal{L} \longmapsto \mathcal{L}_S$$

induces a bijection between the sets of strict localizations of \mathcal{C} and \mathcal{C}_S .

Note that this theorem *does not* say that any nilpotent deformation of a localization of \mathcal{C}_S is itself a localization of \mathcal{C} . This is in fact false (even under the appropriate flatness hypotheses).

Example 7.2. Consider $S = k$, a field. $R = k[\epsilon]$, $\epsilon^2 = 0$. $\Lambda = k[x, y]$. $\mathcal{C} = \text{Mod}(\Lambda)$, $\mathcal{L} = \text{Mod}(\Lambda_x)$. By Theorem 8.16 below the deformations of \mathcal{C} and \mathcal{L} correspond to the deformations of Λ and Λ_x . These are respectively given by the rings

$$R[x, y]/(yx - xy - f\epsilon) \quad \text{and} \\ R[x, x^{-1}, y]/((yx - xy - g\epsilon))$$

for $f \in k[x, y]$, $g \in k[x, x^{-1}, y]$. Clearly Λ_x has many more deformations than Λ .

The next theorem allows us to recognize those deformations of localizations which are themselves localizations.

Theorem 7.3. *Consider a commutative diagram*

$$(7.1) \quad \begin{array}{ccc} \mathcal{C} & \begin{array}{c} \xrightarrow{a'} \\ \xleftarrow{i'} \end{array} & \mathcal{L} \\ F \uparrow & & \uparrow F' \\ \mathcal{C}_S & \begin{array}{c} \xrightarrow{a} \\ \xleftarrow{i} \end{array} & \mathcal{L}_S \end{array}$$

in which F and F' are flat nilpotent deformations of Grothendieck categories and (a, i) and (a', i') are pairs of adjoint functors. Suppose i' maps injectives onto coflats. If i is a localization then the same holds for i' .

If $u : \mathbf{u} \rightarrow \mathcal{C}$ is an additive functor from a small pre-additive category \mathbf{u} to a Grothendieck category \mathcal{C} , then we have a pair of adjoint functors (a, i) where $i : \mathcal{C} \rightarrow \text{Pr}(\mathbf{u}) : U \mapsto \mathcal{C}(u(-), U)$ and $a : \text{Pr}(\mathbf{u}) \rightarrow \mathcal{C}$ is the unique colimit preserving functor sending $\mathbf{u}(-, A)$ to $u(A)$. We say that u induces a localization if i is a localization.

We will apply Theorem 7.3 in the following setting: assume that $R \rightarrow S$ has nilpotent kernel and consider a commutative diagram

$$(7.2) \quad \begin{array}{ccc} \mathbf{u} & \xrightarrow{u} & \mathcal{C} \\ f \downarrow & & \downarrow (S \otimes_R -) \\ \mathbf{v} & \xrightarrow{v} & \mathcal{D} \end{array}$$

where the categories and functors are as follows:

- (1) \mathbf{u}, \mathbf{v} are respectively small R and S -linear flat categories;
- (2) the functor f is a nilpotent linear deformation;
- (3) \mathcal{C}, \mathcal{D} are respectively R and S -linear flat Grothendieck categories;
- (4) u, v are respectively R and S -linear additive functors;
- (5) the functor $S \otimes_R -$ is the left adjoint to a deformation $\mathcal{D} \rightarrow \mathcal{C}$;
- (6) the images of u, v consist of flat objects.

Proposition 7.4. *If v induces a localization $\mathcal{D} \rightarrow \text{Pr}(\mathbf{v})$, then u induces a localization $\mathcal{C} \rightarrow \text{Pr}(\mathbf{u})$.*

7.2. Proofs. Let \mathcal{C} be an abelian category. Recall that a full subcategory $\mathcal{S} \subset \mathcal{C}$ closed under subquotients and extensions is called a *Serre subcategory* of \mathcal{C} . If moreover \mathcal{C} is Grothendieck and \mathcal{S} is closed under coproducts, then \mathcal{S} is called a *localizing subcategory* of \mathcal{C} . If \mathcal{S} is localizing then the quotient category \mathcal{C}/\mathcal{S} is again

a Grothendieck category and the composition $\mathcal{S}^\perp \longrightarrow \mathcal{C} \longrightarrow \mathcal{C}/\mathcal{S}$, where \mathcal{S}^\perp is the full subcategory of \mathcal{S} with objects

$$\text{Ob}(\mathcal{S}^\perp) = \{C \in \mathcal{C} \mid \forall S \in \mathcal{S} : \text{Ext}_{\mathcal{C}}^i(S, C) = 0, i = 0, 1\},$$

is an equivalence of categories.

Assume that $i : \mathcal{L} \longrightarrow \mathcal{C}$ is a localization of Grothendieck categories with exact left adjoint a . Then $\ker(a)$ is a localizing subcategory and a induces an equivalence $\mathcal{C}/\ker(a) \longrightarrow \mathcal{L}$.

If i is a strict localization then $\mathcal{L} = \ker(a)^\perp$. We summarize this discussion in the following proposition:

Proposition 7.5. *Consider a Grothendieck category \mathcal{C} . Let \mathfrak{s} denote the set of localizing subcategories of \mathcal{C} and let \mathfrak{l} denote the set of strict localizations of \mathcal{C} . Then*

$$\mathfrak{s} \longrightarrow \mathfrak{l} : \mathcal{S} \longmapsto \mathcal{S}^\perp$$

and

$$\mathfrak{l} \longrightarrow \mathfrak{s} : (\mathcal{L} \xrightarrow{i} \mathcal{C}) \longmapsto \ker(a),$$

where a is a left adjoint of i , are inverse bijections. \square

For a set of objects or a subcategory $\mathcal{S} \subset \mathcal{C}$ we will write $\langle \mathcal{S} \rangle_{\mathcal{C}}$ for the smallest Serre subcategory of \mathcal{C} containing \mathcal{S} .

Now assume that R, S, I have their usual meaning and assume $I^n = 0$. Let \mathcal{C} be an R -linear abelian category.

Proposition 7.6. *Let \mathfrak{s}' denote the set of Serre subcategories of \mathcal{C} and \mathfrak{s} the set of Serre subcategories of \mathcal{C}_S . Then*

$$\mathfrak{s}' \longrightarrow \mathfrak{s} : \mathcal{S}' \longmapsto \mathcal{S}' \cap \mathcal{C}_S$$

and

$$\mathfrak{s} \longrightarrow \mathfrak{s}' : \mathcal{S} \longmapsto \langle \mathcal{S} \rangle_{\mathcal{C}}$$

are inverse bijections. If \mathcal{C} is a Grothendieck category, they restrict to inverse bijections between the respective sets of localizing subcategories.

Proof. Let $\mathcal{S}, \mathcal{S}'$ be objects in \mathfrak{s} and \mathfrak{s}' respectively. Any object C in \mathcal{C} has a finite filtration

$$0 = I^n C \subseteq I^{n-1} C \subseteq \dots \subseteq I^2 C \subseteq IC \subseteq I^0 C = C$$

Write $\text{gr } C \in \mathcal{C}_S$ for its associated graded object. The formation of $\text{gr } C$ is compatible with coproducts.

For $\mathcal{S} \in \mathfrak{s}$ let $\overline{\mathcal{S}}$ be the full subcategory of \mathcal{C} whose objects are given by

$$\text{Ob}(\overline{\mathcal{S}}) = \{C \in \mathcal{C} \mid \text{gr } C \in \mathcal{S}\}$$

It is easy to see that $\overline{\mathcal{S}}$ is a Serre subcategory of \mathcal{C} which is localizing if \mathcal{S} is.

Clearly $\mathcal{S} \subset \overline{\mathcal{S}} \subset \langle \mathcal{S} \rangle_{\mathcal{C}}$ and since $\overline{\mathcal{S}}$ is Serre, we deduce $\langle \mathcal{S} \rangle_{\mathcal{C}} = \overline{\mathcal{S}}$. This immediately implies

$$\langle \mathcal{S} \rangle_{\mathcal{C}} \cap \mathcal{C}_S = \mathcal{S}$$

Clearly if $\mathcal{S}' \in \mathfrak{s}'$ then $\mathcal{S}' \cap \mathcal{C}_S \in \mathfrak{s}$, and since colimits in \mathcal{C} and \mathcal{C}_S are computed in the same way (Proposition 4.2), $\mathcal{S}' \cap \mathcal{C}_S$ is localizing if \mathcal{S}' is. We certainly have

$$\langle \mathcal{S}' \cap \mathcal{C}_S \rangle_{\mathcal{C}} \subset \mathcal{S}'$$

However if $C \in \mathcal{S}'$ then $\text{gr } C \in \mathcal{S}' \cap \mathcal{C}_S$ and hence $C \in \langle \mathcal{S}' \cap \mathcal{C}_S \rangle_{\mathcal{C}}$ by the earlier discussion. Thus in fact $\langle \mathcal{S}' \cap \mathcal{C}_S \rangle_{\mathcal{C}} = \mathcal{S}'$ and we are done. \square

Proof of Theorem 7.1. Let $i : \mathcal{L} \rightarrow \mathcal{C}$ be a strict localization and let $a : \mathcal{C} \rightarrow \mathcal{L}$ be the exact left adjoint to i . Consider the pair of functors (i_S, a_S) between \mathcal{L}_S and \mathcal{C}_S . It is clear that i_S is still fully faithful and that a_S is an exact left adjoint to i_S . Thus $i_S : \mathcal{L}_S \rightarrow \mathcal{C}_S$ is a (strict) localization.

Let \mathfrak{l}' denote the set of strict localizations of \mathcal{C} and \mathfrak{l} the set of strict localizations of \mathcal{C}_S . Propositions 7.5 and 7.6 furnish us with bijections

$$\mathfrak{l}' \longrightarrow \mathfrak{s}' \longrightarrow \mathfrak{s} \longrightarrow \mathfrak{l}$$

whose composition sends \mathcal{L} to $(\ker(a) \cap \mathcal{C}_S)^\perp = \ker(a_S)^\perp = \mathcal{L}_S$, finishing the proof. \square

The proof of Theorem 7.3 is based on the following observation:

Proposition 7.7. *Consider a functor $i : \mathcal{L} \rightarrow \mathcal{C}$ between Grothendieck categories and suppose i has a left adjoint a . The following are equivalent:*

- (1) i is a localization;
- (2) i preserves injectives and $i : \text{Inj}(\mathcal{L}) \rightarrow \text{Inj}(\mathcal{C})$ is fully faithful.

Proof. Obviously (1) implies (2). Suppose (2) holds. It is easily seen that i is fully faithful. The exactness of a follows since exactness of a sequence can be tested by considering Hom 's into all injectives. \square

Proof of Theorem 7.3. We use Proposition 7.7. First, let us show that i' preserves injectives. Consider $E \in \text{Inj}(\mathcal{L})$. To prove that $i'(E)$ is injective in \mathcal{C} , it suffices that for all $C \in \mathcal{C}_S$ we have $\text{Ext}_{\mathcal{C}}^1(C, i'(E)) = \text{Ext}_{\mathcal{C}_S}^1(C, \text{Hom}_R(S, i'(E))) = 0$ where the first equality follows from Proposition 4.7.

Looking at right adjoints in (7.1) we obtain a commutative diagram

$$\begin{array}{ccc} \mathcal{L} & \xrightarrow{i'} & \mathcal{C} \\ \text{Hom}_R(S, -) \downarrow & & \downarrow \text{Hom}_R(S, -) \\ \mathcal{L}_S & \xrightarrow{i} & \mathcal{C}_S. \end{array}$$

The desired result follows from the fact that $\text{Hom}_R(S, i'(E)) = i(\text{Hom}_R(S, E))$ is injective in \mathcal{C}_S . Next, we are to prove that $\eta : \mathcal{L}(I, I') \rightarrow \mathcal{C}(i'(I), i'(I'))$ is an isomorphism for injectives I and I' . Since $\mathcal{L}(I, I')$ and $\mathcal{C}(i'(I), i'(I'))$ are flat in $\text{Mod}(R)$, using the 5-lemma it suffices that $S \otimes_R \eta$ is an isomorphism. But $S \otimes_R \eta$ is isomorphic to $\mathcal{L}_S(\text{Hom}_R(S, I), \text{Hom}_R(S, I')) \rightarrow \mathcal{C}_S(i(\text{Hom}_R(S, I)), i(\text{Hom}_R(S, I')))$, which proves our assertion. \square

Proof of Proposition 7.4. We apply Theorem 7.3. We have a diagram

$$(7.3) \quad \begin{array}{ccc} \text{Pr}(\mathbf{u}) & \xrightleftharpoons[a']{a'} & \mathcal{C} \\ F \uparrow & & \uparrow F' \\ \text{Pr}(\mathbf{v}) & \xrightleftharpoons[i]{a} & \mathcal{D} \end{array}$$

where (a, i) , (a', i') are the pairs of adjoint functors associated to v and u ; F' is the deformation we started with and F is the abelian deformation associated to the linear deformation $f^{\text{op}} : \mathbf{u}^{\text{op}} \rightarrow \mathbf{v}^{\text{op}}$ (see Proposition 5.4).

We claim that (7.3) is commutative. We first consider the “ i -square”. Starting with $D \in \mathcal{D}$ we have

$$\begin{aligned} (F \circ i)(D) &= \mathcal{D}((v \circ f)(-), D) \\ &= \mathcal{D}(S \otimes_R u(-), D) \\ &= \mathcal{C}(u(-), F'(D)) \\ &= (i' \circ F')(D). \end{aligned}$$

Since F, F' are fully faithful any R -linear functor G between $\text{Pr}(\mathbf{u})$ and \mathcal{C} (in both directions) restricts to exactly one functor between $\text{Pr}(\mathbf{v})$ and \mathcal{D} (up to natural isomorphism) necessarily given by G_S . A pair of adjoint functors (G, H) restricts to a pair of adjoint functors (G_S, H_S) .

Thus we obtain a pair of adjoint functors (a'_S, i'_S) between $\text{Pr}(\mathbf{u})$ and \mathcal{C} and furthermore $i = i'_S$. Since a is the left adjoint to i we obtain $a = a'_S$ (up to natural isomorphism). This implies the commutivity of the “ a -square” in (7.3).

The remaining hypothesis of Theorem 7.3 we need to check is that i' sends injectives to coflats. If $E \in \mathcal{C}$ is injective then it is coflat by Proposition 3.4. Since the objects of \mathbf{u} are mapped to flat objects by u this implies that $\mathcal{C}(u(-), E)$ takes on coflat values by Proposition 2.9(7). Hence by Proposition 2.10 $i'(E)$ is coflat. \square

8. EQUIVALENT DEFORMATION PROBLEMS

In this section we obtain several “equivalences of deformation problems”. We start by formalizing what we mean by this.

8.1. Deformation pseudo functors. In this section we need to be careful about our choices of universe. Therefore we make them temporarily explicit in our notations.

Let \mathcal{U} be a universe. We will denote by $\mathcal{U}\text{-Rng}^0$ the category with as objects coherent commutative \mathcal{U} -rings and as morphisms surjective ring maps with a finitely generated, nilpotent kernel. For a fixed coherent ring $S \in \mathcal{U}$, we consider the category $\mathcal{U}\text{-Rng}^0/S$.

Fix some other universe \mathcal{W} . A deformation pseudo functor is by definition a pseudo functor

$$D : \mathcal{U}\text{-Rng}^0/S \longrightarrow \mathcal{W}\text{-Gd}$$

An *equivalence* of deformation pseudo functors $D_1, D_2 : \mathcal{U}\text{-Rng}^0/S \longrightarrow \mathcal{W}\text{-Gd}$ is a pseudonatural transformation $\mu : D_1 \longrightarrow D_2$ for which every $D_1(R) \longrightarrow D_2(R)$ is an equivalence of categories. It is easy to see that this defines an equivalence relation on deformation pseudo functors.

The dependence of our notations on the universes \mathcal{U} and \mathcal{W} is a nuisance, but the deformation pseudo functors we will consider below will be stable under enlarging \mathcal{U} and \mathcal{W} in a suitable sense. The following proposition is a first step in this direction.

Proposition 8.1. *Let $\mathcal{U} \subset \mathcal{U}'$ be universes. Then the obvious functor*

$$\mathcal{U}\text{-Rng}^0/S \longrightarrow \mathcal{U}'\text{-Rng}^0/S$$

is an equivalence of categories. Consequently, two deformation pseudofunctors $D_1, D_2 : \mathcal{U}'\text{-Rng}^0/S \longrightarrow \mathcal{W}\text{-Gd}$ are equivalent if and only if their restrictions to $\mathcal{U}\text{-Rng}^0/S$ are.

Proof. We only need to show that it is essentially surjective, and this readily follows from Proposition 6.5. \square

8.2. Abelian and linear deformations.

Definition 8.2. (1) Consider an S -linear category \mathfrak{b} . An *equivalence of deformations* from $f_1 : \mathfrak{a}_1 \rightarrow \mathfrak{b}$ to $f_2 : \mathfrak{a}_2 \rightarrow \mathfrak{b}$ is an equivalence of R -linear categories $\varphi : \mathfrak{a}_2 \rightarrow \mathfrak{a}_1$ such that $f_1 \circ \varphi$ is naturally isomorphic to f_2 . If φ is an isomorphism with $f_1 \circ \varphi = f_2$, it is called an *isomorphism of deformations*.

(2) Consider an abelian S -linear category \mathcal{D} . An *equivalence of deformations* from $F_1 : \mathcal{D} \rightarrow \mathcal{C}_1$ to $F_2 : \mathcal{D} \rightarrow \mathcal{C}_2$ is an equivalence of R -linear categories $\Phi : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ such that $\Phi \circ F_1$ is naturally isomorphic to F_2 . If Φ is an isomorphism with $\Phi \circ F_1 = F_2$, it is called an *isomorphism of deformations*.

We consider the following groupoids for $R \in \mathcal{U}\text{-Rng}^0/S$ (where $S \in \mathcal{U}$).

- (1) For a flat S -linear \mathcal{U} -category \mathfrak{b} and a universe \mathcal{V} such that \mathfrak{b} is essentially \mathcal{V} -small and $\mathcal{U} \in \mathcal{V}$, we consider the groupoid $\mathcal{V}\text{-def}_{\mathfrak{b}}(R)$. The objects of $\mathcal{V}\text{-def}_{\mathfrak{b}}(R)$ are flat R -deformations of \mathfrak{b} which are elements of \mathcal{V} . Its morphisms are equivalences of deformations modulo natural isomorphism of functors.
- (2) For a flat S -linear abelian \mathcal{U} -category \mathcal{D} and a universe \mathcal{V} such that \mathcal{D} is essentially \mathcal{V} -small and $\mathcal{U} \in \mathcal{V}$, we consider the groupoid $\mathcal{V}\text{-Def}_{\mathcal{D}}(R)$. The objects of $\mathcal{V}\text{-Def}_{\mathcal{D}}(R)$ are the flat R -deformations of \mathcal{D} which are elements of \mathcal{V} . Its morphisms are equivalences of deformations modulo natural isomorphism of functors.

Clearly $\text{Ob}(\mathcal{V}\text{-def}_{\mathfrak{b}}(R)) \subset \mathcal{V}$ and $\text{Ob}(\mathcal{V}\text{-Def}_{\mathcal{D}}(R)) \subset \mathcal{V}$. Hence if we take a universe \mathcal{W} with $\mathcal{V} \in \mathcal{W}$, this yields deformation pseudo functors

$$\mathcal{V}\text{-def}_{\mathfrak{b}}, \mathcal{V}\text{-Def}_{\mathcal{D}} : \mathcal{U}\text{-Rng}^0/S \rightarrow \mathcal{W}\text{-Gd}$$

The universe \mathcal{W} is a purely technical device to make sure $\mathcal{V}\text{-def}_{\mathfrak{b}}$, $\mathcal{V}\text{-Def}_{\mathcal{D}}$ take their values in a category. Obviously, the equivalence or non equivalence of two deformation pseudofunctors $D_1, D_2 : \mathcal{U}\text{-Rng}^0/S \rightarrow \mathcal{W}\text{-Gd}$ is preserved under changing from \mathcal{W} to \mathcal{W}' for some $\mathcal{W} \subset \mathcal{W}'$.

The following proposition shows that the choice of the universe \mathcal{V} is harmless as well. Consider universes $\mathcal{V} \subset \mathcal{V}'$ with $\mathcal{V}' \in \mathcal{W}$.

Proposition 8.3. *Let \mathfrak{b} be a flat S -linear \mathcal{U} -category and let \mathcal{D} be a flat S -linear abelian \mathcal{U} -category. Then the obvious pseudonatural transformations*

$$\begin{aligned} \mathcal{V}\text{-def}_{\mathfrak{b}} &\longrightarrow \mathcal{V}'\text{-def}_{\mathfrak{b}} \\ \mathcal{V}\text{-Def}_{\mathcal{D}} &\longrightarrow \mathcal{V}'\text{-Def}_{\mathcal{D}} \end{aligned}$$

are equivalences of deformation pseudo functors.

Proof. This follows from Lemmas 6.7, 6.9. \square

8.3. Small skeletons. From now on, we will simply write $\text{def}_{\mathfrak{b}}$ and $\text{Def}_{\mathcal{D}}$ to denote the functors $\mathcal{V}\text{-def}_{\mathfrak{b}}$ and $\mathcal{V}\text{-Def}_{\mathcal{D}}$ for some \mathcal{U}, \mathcal{V} and \mathcal{W} as above. Below we fix the universe \mathcal{U} . We assume that all rings are in \mathcal{U} and write Rng^0 for $\mathcal{U}\text{-Rng}^0$. The notions of small and essentially small are as usual with respect to the fixed universe \mathcal{U} . The same holds for the notion of a Grothendieck category.

We prove the following results.

Theorem 8.4. *Assume that \mathfrak{b} is an essentially small flat S -linear category. Then $\text{def}_{\mathfrak{b}}(R)$ has a small skeleton.*

Proof. Since \mathfrak{b} is essentially small, there is an infinite small cardinal κ with $|\text{Sk}(\mathfrak{b})| \leq \kappa$ and $|\mathfrak{b}(B, B')| \leq \kappa$ for all $B, B' \in \mathfrak{b}$. By Lemma 6.6, the same bound holds for every $\mathfrak{a} \in \text{def}_{\mathfrak{b}}(R)$. It is easily seen that up to equivalence, the number of R -linear categories \mathfrak{a} satisfying this bound is small. Furthermore, for a given \mathfrak{a} , up to natural isomorphism, the number of equivalences $S \otimes_R \mathfrak{a} \rightarrow \mathfrak{b}$ is small (see Lemma 2.3). \square

Theorem 8.5. *Assume that \mathcal{D} is a flat S -linear abelian category which is either essentially small or a Grothendieck category. Then $\text{Def}_{\mathcal{D}}(R)$ has a small skeleton.*

Proof. We may assume that $I^2 = 0$. Assume first that \mathcal{D} is essentially small. Then there is an infinite small cardinal κ with $|\text{Sk}(\mathcal{D})| \leq \kappa$ and $|\mathcal{D}(D, D')| \leq \kappa$ for all $D, D' \in \mathcal{D}$. By Lemma 6.8, the same bound holds for every $\mathcal{C} \in \text{Def}_{\mathcal{D}}(R)$. The proof is then finished like the proof of Theorem 8.4.

Now assume that \mathcal{D} is a Grothendieck category with a generator G . Let κ and \mathcal{D}_{κ} be as in section 6.5. If $\mathcal{D} \rightarrow \mathcal{C}$ is a deformation, the image G' of G in \mathcal{C}_S defines a cardinality on \mathcal{C}_S which allows us to define \mathcal{C}_{κ} as in section 6.5. By Lemma 6.20, we can take an infinite small cardinal λ with $|\text{Sk}(\mathcal{D}_{\kappa})| \leq \lambda$ and $|\mathcal{D}(D, D')| \leq \lambda$ for all $D, D' \in \mathcal{D}_{\kappa}$. By Lemma 6.8, the same bound holds for \mathcal{C}_{κ} for every $\mathcal{C} \in \text{Def}_{\mathcal{D}}(R)$, and by Lemmas 6.23, 6.24, \mathcal{C}_{κ} generates \mathcal{C} . Consequently, using Theorem 6.18, up to equivalence, a deformation $\mathcal{D} \rightarrow \mathcal{C}$ is determined by:

- (1) an R -linear category \mathfrak{g} with $|\text{Ob}(\mathfrak{g})| \leq \lambda$ and $|\mathfrak{g}(G, G')| \leq \lambda$ for all $G, G' \in \mathfrak{g}$;
- (2) an additive topology on \mathfrak{g} ;
- (3) an object $G' \in \mathfrak{g}$ and a map $\mathcal{D}(G, G) \rightarrow \mathfrak{g}(G', G')$.

Clearly, the number of such data is small, which finishes the proof. \square

Remark 8.6. Example 6.17 shows that Theorem 8.5 is trivially false without the flatness requirement on deformations, even for $\mathcal{D} = \text{Mod}(k)$, k a field and $R = k[\epsilon]/(\epsilon^2)$. Indeed every small cardinal κ in Example 6.17 yields a different deformation of \mathcal{D} and the set of small cardinals is not itself small.

8.4. Elementary deformation equivalences. We have the following elementary deformation equivalences.

Proposition 8.7. (1) *For every flat S -linear \mathfrak{b} there is an equivalence*

$$\theta : \text{def}_{\mathfrak{b}} \rightarrow \text{def}_{\mathfrak{b}^{\text{op}}};$$

(2) *an equivalence $F : \mathfrak{b} \rightarrow \mathfrak{b}'$ of flat S -linear categories induces an equivalence*

$$\delta : \text{def}_{\mathfrak{b}} \rightarrow \text{def}_{\mathfrak{b}'};$$

(3) *for every flat abelian \mathcal{D} there is an equivalence*

$$\theta : \text{Def}_{\mathcal{D}} \rightarrow \text{Def}_{\mathcal{D}^{\text{op}}};$$

(4) *an equivalence $F : \mathcal{D} \rightarrow \mathcal{D}'$ of flat abelian categories induces an equivalence*

$$\delta : \text{Def}_{\mathcal{D}} \rightarrow \text{Def}_{\mathcal{D}'}.$$

Proof. The only non-trivial point here is that by Proposition 3.3, the opposite of a flat abelian category is again flat. Note that the corresponding statement for R -linear categories is obviously true. \square

8.5. Ind-objects versus coherent objects. If \mathcal{C}_{es} is essentially small then there is an equivalence $\mathcal{C}_{es} \cong \mathbf{Fp}(\mathbf{Ind}(\mathcal{C}_{es}))$. Similarly for a locally coherent Grothendieck category \mathcal{C} there is an equivalence $\mathbf{Ind}(\mathbf{Fp}(\mathcal{C})) \cong \mathcal{C}$ (see §2.2).

Using Proposition 5.5 and Proposition 5.6, we obtain pseudo natural transformations

$$(8.1) \quad \mathbf{Def}_{\mathcal{D}_{es}} \longrightarrow \mathbf{Def}_{\mathbf{Ind}(\mathcal{D}_{es})}$$

and

$$(8.2) \quad \mathbf{Def}_{\mathcal{D}} \longrightarrow \mathbf{Def}_{\mathbf{Fp}(\mathcal{D})}$$

for an essentially small S -linear category \mathcal{D}_{es} and an S -linear locally coherent Grothendieck category \mathcal{D} . By Proposition 6.9, any $\mathcal{C} \in \mathbf{Def}_{\mathcal{D}_{es}}$ is essentially small, and by Theorem 6.36, any $\mathcal{C} \in \mathbf{Def}_{\mathcal{D}}$ is a locally coherent Grothendieck category. This allows one to prove the following

Theorem 8.8. *(8.1) and (8.2) are equivalences of deformation pseudo functors.* \square

8.6. Lifting deformations of localizations. In this section we prove a general result (Theorem 8.14 below) which will allow us to construct deformation equivalences in several settings afterwards. Unfortunately the formulation of Theorem 8.14 is rather technical so the interested reader may consider reading the subsequent sections first.

Our purpose is to describe a class of functors $v : \mathfrak{v} \longrightarrow \mathcal{D}$, with \mathfrak{v} flat pre-additive and \mathcal{D} flat abelian, for which $\mathbf{def}_{\mathfrak{v}}$ and $\mathbf{Def}_{\mathcal{D}}$ are equivalent.

Consider a diagram

$$\begin{array}{ccc} \mathfrak{u} & & \mathcal{C} \\ f \downarrow & & \downarrow (S \otimes_R -) \\ \mathfrak{v} & \xrightarrow{u} & \mathcal{D} \end{array}$$

in which v is an S -linear functor from a small S -linear category to an S -linear Grothendieck category, $(S \otimes_R -)$ is left adjoint to an abelian R -deformation $F : \mathcal{D} \longrightarrow \mathcal{C}$ and f is a linear R -deformation. In Section 7.1, we have concentrated on lifting properties of v , a lift of v being given. Now we will concentrate on the problem of lifting v . We have the following interpretation:

Proposition 8.9. *The canonical functor*

$$\mathbf{Add}(S)(\mathfrak{v}, \mathcal{D}) \xrightarrow{\cong} \mathbf{Add}(R)(\mathfrak{u}, \mathcal{D}) \xrightarrow{(F \circ \cdot)} \mathbf{Add}(R)(\mathfrak{u}, \mathcal{C})$$

is an abelian R -deformation, its left adjoint being given by

$$\mathbf{Add}(R)(\mathfrak{u}, \mathcal{C}) \xrightarrow{(S \otimes_R \circ \cdot)} \mathbf{Add}(R)(\mathfrak{u}, \mathcal{D}) \xrightarrow{\cong} \mathbf{Add}(S)(\mathfrak{v}, \mathcal{D}).$$

Hence an R -linear $u : \mathfrak{u} \longrightarrow \mathcal{C}$ making the diagram commute is a lift of v along this left adjoint. \square

Proposition 8.10. *Consider an S -linear functor $v : \mathfrak{v} \longrightarrow \mathcal{D}$ from a small S -linear category to a S -linear Grothendieck category such that the objects $v(V)$ are flat. Suppose $\mathbf{Ob}(\mathfrak{v})$ is endowed with a reflexive, transitive relation \mathcal{R} such that*

- (1) (U, V) not in \mathcal{R} implies $\mathfrak{v}(U, V) = 0$;

- (2) (U, V) in \mathcal{R} implies that $v_{(U,V)} : \mathfrak{v}(U, V) \rightarrow \mathcal{D}(v(U), v(V))$ is an isomorphism and that $\text{Ext}_{\mathcal{D}}^i(v(U), M \otimes_S v(V)) = 0$ for $M \in \text{mod}(S)$ and $i = 1, 2$.

Consider a flat nilpotent R -deformation $F : \mathcal{D} \rightarrow \mathcal{C}$. The following hold:

- (1) There is a strict flat left R -deformation $f : \mathfrak{u} \rightarrow \mathfrak{v}$ and an R -linear functor $u : \mathfrak{u} \rightarrow \mathcal{C}$ for which all objects $u(U)$ are flat such that the diagram

$$\begin{array}{ccc} \mathfrak{u} & \xrightarrow{u} & \mathcal{C} \\ f \downarrow & & \downarrow (S \otimes_R -) \\ \mathfrak{v} & \xrightarrow{v} & \mathcal{D} \end{array}$$

commutes up to natural isomorphism.

- (2) If $f' : \mathfrak{u}' \rightarrow \mathfrak{v}$ and $u' : \mathfrak{u}' \rightarrow \mathcal{C}$ are like f and u (but with f' not necessarily strict), there is an equivalence $w : \mathfrak{u}' \rightarrow \mathfrak{u}$ with $f \circ w = f'$ and $v \circ w \cong v'$.

Proof. We will construct u as stated in case $I^2 = 0$. It will be clear from the proof and from Proposition 6.13 that the R -linear functor $u : \mathfrak{u} \rightarrow \mathcal{C}$ satisfies the same properties as v , which then finishes the proof in the general case.

For (1), consider a diagram

$$\begin{array}{ccc} & & \mathcal{C} \\ & & \downarrow (S \otimes_R -) \\ \mathfrak{v} & \xrightarrow{v} & \mathcal{D} \end{array}$$

as stated. By Theorem 6.12, $v(V)$ has a flat lift $D_V \rightarrow v(V)$ along the functor $(S \otimes_R -)$. Define \mathfrak{u} by $\text{Ob}(\mathfrak{u}) = \text{Ob}(\mathfrak{v})$ and $\mathfrak{u}(U, V) = \mathcal{C}(D_U, D_V)$ if (U, V) is in \mathcal{R} and $\mathfrak{u}(U, V) = 0$ else. Since \mathcal{R} is transitive, there is an obvious composition on \mathfrak{u} making \mathfrak{u} into an R -linear category with an R -linear functor $u : \mathfrak{u} \rightarrow \mathcal{C}$. There is also an obvious functor $f : \mathfrak{u} \rightarrow \mathfrak{v}$. For (U, V) not in \mathcal{R} , $f_{(U,V)} : 0 = \mathfrak{u}(U, V) \rightarrow \mathfrak{v}(U, V) = 0$ is trivially a deformation of modules. For (U, V) in \mathcal{R} , $f_{(U,V)}$ is isomorphic to $(S \otimes_R -)_{(U,V)} : \mathcal{C}(D_U, D_V) \rightarrow \mathcal{D}(S \otimes_R D_U, S \otimes_R D_V)$. It follows from Proposition 2.9(8) that $\mathfrak{v}(U, V)$ is flat and $f_{(U,V)}$ is a deformation of modules. Hence f is a flat left R -deformation, and the diagram commutes up to natural isomorphism.

For (2), consider f' and u' as stated. Since every $u'(W)$ is flat and we have natural isomorphisms $S \otimes_R u'(W) \rightarrow v(f'(W))$, Theorem 6.12 provides \mathcal{D} -isomorphisms $u'(W) \rightarrow D_{f'(W)}$ that allow the definition of a functor $w : \mathfrak{u}' \rightarrow \mathfrak{u}$. w is essentially surjective since f' is a deformation. For $(f'(U), f'(V))$ not in \mathcal{R} , since $f'_{(U,V)} : \mathfrak{u}'(U, V) \rightarrow \mathfrak{v}(f'(U), f'(V)) = 0$ is a nilpotent deformation of modules, it follows that $\mathfrak{u}'(U, V) = 0$. Next, suppose $(f'(U), f'(V))$ is in \mathcal{R} . We have a commutative diagram

$$\begin{array}{ccc} \mathfrak{u}'(U, V) & \xrightarrow{u'_{(U,V)}} & \mathcal{C}(u'(U), u'(V)) \\ f'_{(U,V)} \downarrow & & \downarrow \\ \mathfrak{v}(f'(U), f'(V)) & \xrightarrow{v_{(f'(U), f'(V))}} & \mathcal{D}(v f'(U), v f'(V)) \end{array}$$

in which the right arrow is isomorphic to $(S \otimes_R -)_{(u'(U), u'(V))}$ and hence both vertical arrows are flat deformations of modules and $v_{(f'(U), f'(V))}$ is an isomorphism. Since

$I \otimes_R u'_{(U,V)} \cong I \otimes_S v_{(f'(U), f'(V))}$, it follows from the 5-lemma that $u'_{(U,V)}$ and hence also $w_{(U,V)}$ is an isomorphism. This proves that w is fully faithful. \square

Proposition 8.11. *For $v : \mathfrak{v} \longrightarrow \mathcal{D}$ as in Proposition 8.10 and $R \in \text{Rng}^0/S$, there is a functor*

$$\kappa_R : \text{Def}_{\mathcal{D}}(R) \longrightarrow \text{def}_{\mathfrak{v}}(R).$$

Proof. For $F : \mathcal{D} \longrightarrow \mathcal{C}$ in $\text{Def}_{\mathcal{D}}(R)$, we can define $\kappa_R(F)$ as $f : \mathfrak{u} \longrightarrow \mathfrak{v}$ of Proposition 8.10(1). Proposition 8.10(2) can be used to define κ_R on morphisms. \square

Proposition 8.12. *For a localization of S -linear abelian categories*

$$\mathcal{D} \begin{array}{c} \xrightarrow{a} \\ \xleftarrow{i} \end{array} \mathcal{K}$$

and $R \in \text{Rng}^0/S$, there is a functor

$$\lambda_R : \text{Def}_{\mathcal{D}}(R) \longrightarrow \text{Def}_{\mathcal{K}}(R).$$

Proof. Since every localization factors as an equivalence followed by a strict localization, this is a consequence of Theorem 7.1 and Lemma 8.13 below applied to the functor a . \square

Lemma 8.13. *Consider an R -linear exact functor $F : \mathcal{C} \longrightarrow \mathcal{D}$ that is essentially surjective. If \mathcal{C} is flat over R , the same holds for \mathcal{D} .*

Proof. For $D \in \mathcal{D}$ and $X \in \text{mod}(R)$, take an isomorphism $d : D \longrightarrow F(C)$ for a C in \mathcal{C} . Take a \mathcal{C} -monomorphism $c : C \longrightarrow C'$ with $\text{Ext}_{\mathcal{C},R}^1(X, c) = 0$. Then $F(c) \circ d$ is a \mathcal{D} -monomorphism and by Proposition 2.7, $\text{Ext}_{\mathcal{D},R}^1(X, F(c)) = F(\text{Ext}_{\mathcal{C},R}^1(X, c)) = 0$. \square

The construction of an abelian deformation of $\text{Mod}(\mathfrak{b})$ from an R -linear deformation of an essentially small category \mathfrak{b} yields a pseudonatural transformation

$$\mu : \text{def}_{\mathfrak{b}} \longrightarrow \text{Def}_{\text{Mod}(\mathfrak{b})}.$$

Theorem 8.14. *Let $v : \mathfrak{v} \longrightarrow \mathcal{D}$ be as in Proposition 8.10 and suppose v induces a localization*

$$\text{Pr}(\mathfrak{v}) \begin{array}{c} \xrightarrow{a} \\ \xleftarrow{i} \end{array} \mathcal{C}.$$

The functors

$$\text{def}_{\mathfrak{v}}(R) \xrightarrow{\theta_R} \text{def}_{\mathfrak{v} \circ \text{Pr}}(R) \xrightarrow{\mu_R} \text{Def}_{\text{Pr}(\mathfrak{v})}(R) \xrightarrow{\lambda_R} \text{Def}_{\mathcal{D}}(R)$$

and

$$\text{Def}_{\mathcal{D}}(R) \xrightarrow{\kappa_R} \text{def}_{\mathfrak{v}}(R)$$

are inverse equivalences, where λ_R is as in Corollary 8.12, κ_R as in Corollary 8.11, θ_R as in Proposition 8.7 and μ_R as above.

Proof. First, we indicate a natural isomorphism $\lambda_R \circ \mu_R \circ \theta_R \circ \kappa_R \cong 1_{\text{Def}_{\mathcal{D}}(R)}$. For $F : \mathcal{D} \longrightarrow \mathcal{C}$ in $\text{Def}_{\mathcal{D}}(R)$, its image F' under $\mu_R \circ \theta_R \circ \kappa_R$ fits into a diagram

$$\begin{array}{ccc} \text{Pr}(\mathbf{u}) & \xleftarrow{i'} & \mathcal{C} \\ F' \uparrow & & \uparrow F \\ \text{Pr}(\mathbf{v}) & \xleftarrow{i} & \mathcal{D} \end{array}$$

that commutes up to natural isomorphism. By Proposition 7.4, i' is a localization. The construction of $\lambda_R(F')$ yields a diagram

$$\begin{array}{ccccc} \text{Pr}(\mathbf{u}) & \xleftarrow{i'_1} & \mathcal{C}' & \xleftarrow{i'_0} & \mathcal{C} \\ \uparrow & & \uparrow & & \uparrow \\ \text{Pr}(\mathbf{u})_S & \xleftarrow{i'_{1S}} & \mathcal{C}'_S & \xleftarrow{\sim} & \mathcal{C}_S \\ \sim \uparrow & & \swarrow \sim & & \uparrow \sim \\ \text{Pr}(\mathbf{v}) & \xleftarrow{i} & \mathcal{D} & & \end{array}$$

in which i'_1 and i'_{1S} are strict localizations and i'_0 constitutes an equivalence of deformations between F and $\lambda_R(F')$.

Next, we indicate a natural isomorphism $\kappa_R \circ \lambda_R \circ \mu_R \circ \theta_R \cong 1_{\text{def}_{\mathbf{v}}(R)}$. Consider $f' : \mathbf{u}' \longrightarrow \mathbf{v}$ in $\text{def}_{\mathbf{v}}(R)$. Its image $F : \mathcal{D} \longrightarrow \mathcal{C}$ under $\lambda_R \circ \mu_R \circ \theta_R$ corresponds to a localization $i' : \mathcal{C} \longrightarrow \text{Pr}(\mathbf{u}')$. The diagram

$$\begin{array}{ccc} \mathbf{u}' & \xrightarrow{u'} & \mathcal{C} \\ f' \downarrow & & \downarrow (S \otimes_R -) \\ \mathbf{v} & \xrightarrow{v} & \mathcal{D} \end{array}$$

in which u' is induced by a left adjoint of i' and $(S \otimes_R -)$ is a left adjoint of F commutes up to natural isomorphism. The functor $w : \mathbf{u}' \longrightarrow \mathbf{u}$ of Proposition 8.10 constitutes an equivalence of deformations between f' and $\kappa_R(F)$. \square

8.7. Deformations of Grothendieck categories with acyclic generators. Next we will formulate some specializations of Theorem 8.14.

Theorem 8.15. *Consider a flat S -linear Grothendieck category \mathcal{D} and a small full subcategory \mathfrak{g} such that $\text{Ob}(\mathfrak{g})$ is a collection of flat generators of \mathcal{D} and such that for G, G' in \mathfrak{g}*

$$\text{Ext}_{\mathcal{D}}^i(G, M \otimes_S G') = 0$$

for $i = 1, 2$ and $M \in \text{mod}(S)$. For $R \in \text{Rng}^0/S$, there is an equivalence

$$\text{def}_{\mathfrak{g}}(R) \cong \text{Def}_{\mathcal{D}}(R).$$

Proof. It is readily seen that the inclusion $\mathfrak{g} \longrightarrow \mathcal{D}$ satisfies the conditions of Theorem 8.14. \square

Theorem 8.16. *For an essentially small flat S -linear category \mathfrak{b} ,*

$$\mu : \text{def}_{\mathfrak{b}} \longrightarrow \text{Def}_{\text{Mod}(\mathfrak{b})}$$

is an equivalence of deformation pseudo functors.

Proof. Since \mathfrak{b}^{op} is a full subcategory of $\text{Mod}(\mathfrak{b})$ satisfying the conditions of Theorem 8.15, the result follows from Theorem 8.14. \square

8.8. Deformations of categories with enough injectives. We have seen in Proposition 6.25 that if \mathcal{C} is an abelian category with enough injectives, there is an equivalence $\mathcal{C} \cong (\text{mod}(\text{Inj}(\mathcal{C})))^{\text{op}}$. We prove the following

Theorem 8.17. *If \mathcal{D} has enough injectives then $\text{Def}_{\mathcal{D}}$ and $\text{def}_{\text{Inj}(\mathcal{D})}$ are equivalent deformation pseudo functors.*

Proof. Since $\text{mod}(\text{Inj}(\mathcal{C})) \cong \mathcal{V}\text{-Fp}(\mathcal{V}\text{-Mod}(\text{Inj}(\mathcal{C})))$ where $\text{Inj}(\mathcal{C}) \in \mathcal{V}$, we obtain a pseudo natural transformation

$$(8.3) \quad \text{def}_{\text{Inj}(\mathcal{C})} \longrightarrow \text{Def}_{(\text{mod}(\text{Inj}(\mathcal{C})))^{\text{op}}},$$

which is actually an equivalence by Theorems 8.16, 8.8 and Proposition 8.7. \square

In Appendix A, we clarify this equivalence a little further using certain preservation properties of flat nilpotent linear deformations.

8.9. Sheaves of modules over a ringed space. We will now give an application of Theorem 8.14 to sheaves of modules over a ringed space.

Let X be a topological space and let \mathcal{O}_X be a sheaf of S -algebra's on X . For an open $U \subset X$, let \mathcal{O}_U be the restriction of \mathcal{O}_X to U and denote by $\text{Mod}(\mathcal{O}_U)$ and $\text{PreMod}(\mathcal{O}_U)$ the (S -linear) categories of sheaves and presheaves of \mathcal{O}_U -modules. For a basis \mathcal{B} of the topology, denote by $\mathcal{O}_{\mathcal{B}}$ the restriction of \mathcal{O}_X to \mathcal{B} and by $\text{PreMod}(\mathcal{O}_{\mathcal{B}})$ the category of presheaves of $\mathcal{O}_{\mathcal{B}}$ -modules.

Theorem 8.18. *Let X be a topological space and let \mathcal{O}_X be a sheaf of flat S -algebra's on X as above. Suppose the topology has a basis \mathcal{B} such that for every $U \in \mathcal{B}$, the sheaf cohomology*

$$H^i(U, M \otimes_S \mathcal{O}_U) = 0$$

for $i = 1, 2$ and $M \in \text{mod}(S)$. For $R \in \text{Rng}^0/S$, there is an equivalence

$$\text{Def}_{\text{PreMod}(\mathcal{O}_{\mathcal{B}})}(R) \cong \text{Def}_{\text{Mod}(\mathcal{O}_X)}(R).$$

Proof. Let $a : \text{PreMod}(\mathcal{O}_X) \longrightarrow \text{Mod}(\mathcal{O}_X)$ be the exact sheafication functor left adjoint to inclusion. For an open $U \subset X$, let $j_U : U \longrightarrow X$ be the inclusion map and let $P_U = j_{U,!}^p \mathcal{O}_U$ and $S_U = j_{U,!} \mathcal{O}_U = a(j_{U,!}^p \mathcal{O}_U)$ be the extensions by zero of \mathcal{O}_U in the categories of presheaves and sheaves. Also, let P_U^b denote the restriction of P_U to \mathcal{B} . For any basis \mathcal{B} , for $U \in \mathcal{B}$ and $F \in \text{PreMod}(\mathcal{O}_{\mathcal{B}})$, we have $\text{Hom}(P_U^b, F) = F(U)$. For U and V in \mathcal{B} , we obtain

$$(8.4) \quad \text{Hom}(P_U^b, P_V^b) = \text{Hom}(P_U, P_V) = \begin{cases} \mathcal{O}_X(U) & \text{if } U \subset V \\ 0 & \text{otherwise} \end{cases}$$

and

$$(8.5) \quad \text{Hom}(S_U, S_V) = \{f \in \mathcal{O}_X(U) \mid f \text{ is zero on a neighbourhood of } U \setminus V\}.$$

Thus in particular if $U \subset V$,

$$(8.6) \quad \text{Hom}(P_U, P_V) = \text{Hom}(S_U, S_V).$$

Let \mathfrak{u} be the full subcategory of $\text{PreMod}(\mathcal{O}_X)$ spanned by the objects P_U for $U \in \mathcal{B}$. By (8.4), \mathfrak{u} is isomorphic to the full subcategory of $\text{PreMod}(\mathcal{O}_{\mathcal{B}})$ spanned by

the objects P_U^b for $U \in \mathcal{B}$. Since these objects are a family of finitely generated projective generators of $\text{PreMod}(\mathcal{O}_{\mathcal{B}})$, we deduce that

$$\text{PreMod}(\mathcal{O}_{\mathcal{B}}) \cong \text{Pr}(\mathfrak{u})$$

hence it suffices to prove that

$$\text{def}_{\mathfrak{u}}(R) \cong \text{Def}_{\text{Mod}(\mathcal{O}_X)}(R).$$

Let

$$u : \mathfrak{u} \longrightarrow \text{Mod}(\mathcal{O}_X)$$

be the restriction of a . We will show that u satisfies the conditions of Proposition 8.14. It is easily seen that u induces a localization (see for example [15]), so it remains to verify the conditions stated in Proposition 8.10. If we say that (P_U, P_V) is in \mathcal{R} if $U \subset V$, the result follows from (8.4), (8.6) and the following computation for $U \subset V$

$$\begin{aligned} \text{Ext}_{\text{Mod}(\mathcal{O}_X)}^i(S_U, M \otimes_S S_V) &= \text{Ext}_{\text{Mod}(\mathcal{O}_U)}^i(\mathcal{O}_U, (M \otimes_S S_V)|_U) \\ &= \text{Ext}_{\text{Mod}(\mathcal{O}_U)}^i(\mathcal{O}_U, M \otimes_S \mathcal{O}_U) \\ &= H^i(U, M \otimes_S \mathcal{O}_U) \end{aligned}$$

where we have used that $j_{U,!}$ is exact and hence its right adjoint restriction functor $(-)_U$ preserves injectives. \square

APPENDIX A.

In this Appendix we indicate a direct proof of Corollary 8.17. This proof makes use of some preservation properties of flat nilpotent linear deformations that may be of independent interest.

A.1. Flat nilpotent linear deformations. let $F : \mathfrak{a} \longrightarrow \mathfrak{b}$ be a flat nilpotent R -deformation of an S -linear category \mathfrak{b} . In this section, we will lift some properties of \mathfrak{b} to \mathfrak{a} . The following is well known in the ring case:

Proposition A.1. *If $f : A_1 \longrightarrow A_2$ in \mathfrak{a} is such that $F(f)$ is an isomorphism in \mathfrak{b} , then f is an isomorphism in \mathfrak{a} .*

Proof. It suffices to consider $\mathfrak{a} \longrightarrow S \otimes_R \mathfrak{a}$ nilpotent of order 2. Suppose $f : A_1 \longrightarrow A_2$ in \mathfrak{a} is an isomorphism in \mathfrak{b} . Take an \mathfrak{a} -map $g : A_2 \longrightarrow A_1$ such that $g \circ f = 1_{A_1}$ and $f \circ g = 1_{A_2}$ in $S \otimes_R \mathfrak{a}$. From the exact rows $0 \longrightarrow I\mathfrak{a}(A, A') \longrightarrow \mathfrak{a}(A, A') \longrightarrow (S \otimes_R \mathfrak{a})(A, A') \longrightarrow 0$ and from $I^2 = 0$ we deduce that

$$(g \circ f - 1_{A_1})^2 = 0$$

and

$$(f \circ g - 1_{A_2})^2 = 0.$$

These equations can be rewritten as

$$(2g - g \circ f \circ g) \circ f = 1_{A_1}$$

and

$$f \circ (2g - g \circ f \circ g) = 1_{A_2},$$

which proves our assertion. \square

Proposition A.2. *If $Z \in \mathfrak{a}$ is such that $F(Z)$ is a zero-object in \mathfrak{b} , then Z is a zero-object in \mathfrak{a} .*

Proof. It suffices to consider $\mathfrak{a} \longrightarrow S \otimes_R \mathfrak{a}$. Suppose Z is a zero-object in $S \otimes_R \mathfrak{a}$. So $(S \otimes_R \mathfrak{a})(A, Z) = 0$ for all \mathfrak{a} -objects A . But then by Nakayama $\mathfrak{a}(A, Z) = 0$ for all A , meaning that Z is a zero-object in \mathfrak{a} . \square

Definition A.3. In a category \mathcal{C} , an *idempotent* is a map e with $e \circ e = e$. An idempotent e *splits* if there exist maps r, s with $e = s \circ r$ and $r \circ s = 1$. A category in which all idempotents split is called *Karoubian*.

Remark A.4. The splitting of an idempotent e is equivalent to the existence of the equalizer of e and 1 and to the existence of the coequalizer of e and 1 . Thus an abelian category is Karoubian. If \mathcal{C} is an abelian category with enough injectives, $\text{Inj}(\mathcal{C})$ is Karoubian too since a retract of an injective is injective.

Proposition A.5. *If \mathfrak{b} is Karoubian, the same holds for \mathfrak{a} .*

Proof. It suffices to consider $\mathfrak{a} \longrightarrow S \otimes_R \mathfrak{a}$ nilpotent of order 2. Let $e : A \longrightarrow A$ be an idempotent in \mathfrak{a} and take maps $s : B \longrightarrow A, r : A \longrightarrow B$ in \mathfrak{a} such that $s \circ r = e$ and $r \circ s = 1$ in $S \otimes_R \mathfrak{a}$. From $s \circ r - e \in I\mathfrak{a}(A, A)$ and $r \circ s - 1 \in I\mathfrak{a}(B, B)$ we obtain the following equations in \mathfrak{a} :

- (1) $(s \circ r - e)^2 = 0$;
- (2) $(r \circ s - 1)^2 = 0$;
- (3) $(r \circ s - 1) \circ r \circ (s \circ r - e) = 0$;
- (4) $(s \circ r - e) \circ s \circ (r \circ s - 1) = 0$.

It follows from (1) that

$$(s + e \circ s - s \circ r \circ s) \circ (r \circ e) = e$$

and from (2) that

$$2(r \circ s) - r \circ s \circ r \circ s = 1.$$

Combining (1) and (3) gives us

$$r \circ e \circ s \circ r = r \circ s \circ r,$$

and combining this with (2) and (4) gives

$$r \circ e \circ s = r \circ s.$$

We can now compute that $(r \circ e) \circ (s + e \circ s - s \circ r \circ s) = 2(r \circ s) - r \circ s \circ r \circ s = 1$. So we have shown that $(s + e \circ s - s \circ r \circ s) : B \longrightarrow A$ and $(r \circ e) : A \longrightarrow B$ give a splitting of e in \mathfrak{a} . \square

Proposition A.6. *If $A, B, C, s_1 : A \longrightarrow C$ and $s_2 : B \longrightarrow C$ in \mathfrak{a} are such that $(F(C), F(s_1), F(s_2))$ is a coproduct of $F(A)$ and $F(B)$ in \mathfrak{b} , then (C, s_1, s_2) is a coproduct of A and B in \mathfrak{a} .*

Proof. It suffices to consider $\mathfrak{a} \longrightarrow S \otimes_R \mathfrak{a}$ nilpotent of order 2. Take maps $p_1 : C \longrightarrow A$ and $p_2 : C \longrightarrow B$ in \mathfrak{a} such that (C, s_1, s_2, p_1, p_2) is a biproduct of A and B in $S \otimes_R \mathfrak{a}$. We obtain the following equations in \mathfrak{a} :

- (1) $(p_1 \circ s_1 - 1)^2 = 0$;
- (2) $(p_1 \circ s_2) \circ (p_2 \circ s_1) = 0$;
- (3) $(p_1 \circ s_1 - 1) \circ (p_1 \circ s_2) = 0$;
- (4) $(p_1 \circ s_2) \circ (p_2 \circ s_2 - 1) = 0$;
- (5) $(s_1 \circ p_1 + s_2 \circ p_2 - 1)^2 = 0$.

Put

$$p'_1 = 2p_1 - p_1 \circ s_1 \circ p_1 - p_1 \circ s_2 \circ p_2$$

and

$$p'_2 = 2p_2 - p_2 \circ s_2 \circ p_2 - p_2 \circ s_1 \circ p_1.$$

It follows from (1) and (2) that $p'_1 \circ s_1 = 2p_1 \circ s_1 - (2p_1 \circ s_1 - 1) - 0 = 1$. From (3) and (4) we find that $p'_1 \circ s_2 = 2p_1 \circ s_2 - p_1 \circ s_2 - p_1 \circ s_2 = 0$. Finally, using (5) we obtain that $s_1 \circ p'_1 + s_2 \circ p'_2 = 1$. Combining these results with their symmetric results (changing the roles of A and B), we have shown that $(C, s_1, s_2, p'_1, p'_2)$ is a biproduct of A and B in \mathfrak{a} , which proves our assertion. \square

Proposition A.7. *If \mathfrak{b} is additive, the same holds for \mathfrak{a} .*

Proof. This immediately follows from Proposition A.2 and Proposition A.6. \square

Definition A.8. A pre-additive category \mathfrak{a} is called *coherent* if $\text{mod}(\mathfrak{a})$ is abelian.

Proposition A.9. *If \mathfrak{b} is coherent, the same holds for \mathfrak{a} .*

Proof. We may assume that $\mathfrak{a}, \mathfrak{b}$ are small. Then the result follows from Proposition 5.4 and Theorem 6.36. \square

We mention the following intrinsic characterization of coherence, which we will not explicitly use.

Definition A.10. [14] Consider a map $f : C \rightarrow C'$ in a pre-additive category \mathcal{C} . A *weak cokernel* of f is a map $g : C' \rightarrow C''$ with $g \circ f = 0$ and such that every map h with $h \circ f = 0$ factorizes as $h = h' \circ g$.

In a triangulated category, the cone of a morphism is a weak cokernel.

Proposition A.11. [14, Lemma 1] *An additive category \mathfrak{a} is coherent if and only if \mathfrak{a} has weak cokernels.*

Remark A.12. If \mathcal{C} is an abelian category with enough injectives, there is an equivalence $\mathcal{C} \cong (\text{mod}(\text{Inj}(\mathcal{C})))^{\text{op}}$ (see Proposition 6.25) hence $\text{Inj}(\mathcal{C})$ is coherent. For a map between injectives, we can first take its cokernel and then a mono to an injective to obtain a weak cokernel in $\text{Inj}(\mathcal{C})$.

A.2. Deformations with enough injectives. We have seen in Theorem 8.17 that if \mathcal{C} is an abelian category with enough injectives, there is an equivalence $\text{Def}_{\mathcal{C}} \cong \text{def}_{\text{Inj}(\mathcal{C})}$. We will now give a different approach to this fact. We start by characterizing “categories of injectives of abelian categories with enough injectives”.

Proposition A.13. *For an abelian category \mathcal{C} with enough injectives, $\text{Inj}(\mathcal{C})$ is an additive, coherent, Karoubian category.*

Proof. $\text{Inj}(\mathcal{C})$ is additive since products of injectives are injective and we already noticed in Remark A.4 that $\text{Inj}(\mathcal{C})$ is Karoubian and in Remark A.12 that $\text{Inj}(\mathcal{C})$ is coherent. \square

Proposition A.14. *For an additive, coherent, Karoubian category \mathfrak{a} , the functor*

$$\mathfrak{a}^{\text{op}} \rightarrow \text{mod}(\mathfrak{a}) : A \mapsto (A, -)$$

induces an equivalence of categories

$$\mathfrak{a}^{\text{op}} \rightarrow \mathfrak{p}$$

to the full subcategory \mathfrak{p} of projectives of $\text{mod}(\mathfrak{a})$.

Proof. Take an object P of \mathfrak{p} and consider a presentation

$$\bigoplus_{i=1}^n \mathfrak{a}(A_i, -) \longrightarrow \bigoplus_{j=1}^m \mathfrak{a}(A_j, -) \longrightarrow P \longrightarrow 0.$$

Since P is projective and \mathfrak{a} is additive, P is a retract of the functor $\mathfrak{a}(\bigoplus_{j=1}^m A_j, -)$. But since \mathfrak{a} is Karoubian, it follows that P is itself representable. \square

In other words, for an additive, coherent, Karoubian category \mathfrak{a} , there is an equivalence $\mathfrak{a} \cong \text{Inj}((\text{mod}(\mathfrak{a}))^{\text{op}})$. Combining this with Proposition A.13, we have thus characterized the “categories of injectives of abelian categories with enough injectives” as being precisely the additive, coherent, Karoubian categories.

Let \mathfrak{b} be a coherent S -linear category. Since $\text{mod}(\mathfrak{b}) \cong \mathcal{V} - \text{Fp}(\mathcal{V} - \text{Mod}(\mathfrak{b}))$ where $\mathfrak{b} \in \mathcal{V}$, we obtain a pseudo natural transformation

$$(A.1) \quad \text{def}_{\mathfrak{b}} \longrightarrow \text{Def}_{(\text{mod}(\mathfrak{b}))^{\text{op}}},$$

which is actually an equivalence by Theorem 8.16, Theorem 8.8 and Proposition 8.7. Taking $\mathfrak{b} = \text{Inj}(\mathcal{D})$, this is precisely the equivalence (8.3) of section 8.8. We will now suggest another proof of this equivalence.

Proposition A.15. *Every flat, nilpotent abelian deformation $\mathcal{D} \longrightarrow \mathcal{C}$ with enough injectives induces a flat linear deformation $\text{Hom}_R(S, -) : \text{Inj}(\mathcal{C}) \longrightarrow \text{Inj}(\mathcal{D})$.*

Proof. Since taking injectives preserves equivalence of categories, we may consider $\mathcal{C}_S \longrightarrow \mathcal{C}$. By Proposition 2.9(5) and Proposition 4.3, the map

$$\mathcal{C}(C, E) \longrightarrow \mathcal{C}(\text{Hom}_R(S, C), E) \longrightarrow \mathcal{C}(\text{Hom}_R(S, C), \text{Hom}_R(S, E))$$

induces an isomorphism $S \otimes_R \mathcal{C}(C, E) \longrightarrow \mathcal{C}(\text{Hom}_R(S, C), \text{Hom}_R(S, E))$ when E is injective. It then follows by Corollary 6.15 that $\text{Hom}_R(S, -) : \text{Inj}(\mathcal{C})^S \longrightarrow \text{Inj}(\mathcal{C}_S)$ is an equivalence. The flatness of $\text{Inj}(\mathcal{C})$ follows from 2.9(6). \square

Let \mathcal{D} be an abelian S -linear category with enough injectives. Proposition A.15 yields a pseudo natural transformation

$$(A.2) \quad \text{Def}_{\mathcal{D}} \longrightarrow \text{def}_{\text{Inj}(\mathcal{D})}.$$

By Theorem 6.16, any $\mathcal{C} \in \text{Def}_{\mathcal{D}}$ has enough injectives, and by Propositions A.13, A.7, A.5 and A.9, any $\mathfrak{a} \in \text{def}_{\mathfrak{b}}$ is an additive, coherent, Karoubian category. This allows one to prove the following

Theorem A.16. *(A.1) and (A.2) are equivalences of deformation pseudofunctors.*

APPENDIX B.

In this Appendix, we consider an alternative deformation pseudo functor $\text{def}_{\mathfrak{b}}^{\mathfrak{s}}$ that can be used to study linear deformations of an S -linear category \mathfrak{b} , and we study its relation with $\text{def}_{\mathfrak{b}}$.

For an S -linear category \mathfrak{b} and for $R \in \text{Rng}^0/S$, consider the following groupoid $\text{def}_{\mathfrak{b}}^{\mathfrak{s}}$: the objects of $\text{def}_{\mathfrak{b}}^{\mathfrak{s}}$ are strict R -deformations $\mathfrak{a} \longrightarrow \mathfrak{b}$. The morphisms of $\text{def}_{\mathfrak{b}}^{\mathfrak{s}}$ are isomorphisms of deformations.

There are obvious functors

$$\sigma_R : \text{def}_{\mathfrak{b}}^{\mathfrak{s}}(R) \longrightarrow \text{def}_{\mathfrak{b}}(R).$$

constituting a pseudonatural transformation $\sigma : \text{def}_{\mathfrak{b}}^s \longrightarrow \text{def}_{\mathfrak{b}}$. As every functor between groupoids does, σ_R induces a function

$$\text{Sk}(\sigma_R) : \text{Sk}(\text{def}_{\mathfrak{b}}^s(R)) \longrightarrow \text{Sk}(\text{def}_{\mathfrak{b}}(R)).$$

We will show that for R in Rng^0/S , $\text{Sk}(\sigma_R)$ is a bijection.

Proposition B.1. *For R in Rng^0/S , σ_R is full.*

Proof. Consider two strict left R -deformations $f_1 : \mathfrak{a}_1 \longrightarrow \mathfrak{b}$ and $f_2 : \mathfrak{a}_2 \longrightarrow \mathfrak{b}$ and an equivalence of deformations $\varphi : \mathfrak{a}_1 \longrightarrow \mathfrak{a}_2$. We will construct an isomorphism of deformations $\varphi' : \mathfrak{a}_1 \longrightarrow \mathfrak{a}_2$ that is naturally isomorphic to φ . Consider a natural isomorphism $\eta : f_1 \longrightarrow f_2 \circ \varphi$. For every \mathfrak{a} -map $a : A \longrightarrow A'$, there is a commutative square

$$\begin{array}{ccc} f_1(A) & \xrightarrow{\eta_A} & f_2(\varphi(A)) \\ f_1(a) \downarrow & & \downarrow f_2(\varphi(a)) \\ f_1(A') & \xrightarrow{\eta_{A'}} & f_2(\varphi(A')). \end{array}$$

Since f_2 is a strict deformation, every A in \mathfrak{a}_1 determines a unique object $\varphi'(A)$ in \mathfrak{a}_2 satisfying

$$f_2(\varphi'(A)) = f_1(A).$$

By Proposition A.1, we can lift the \mathfrak{b} -isomorphisms

$$\eta_A : f_2(\varphi'(A)) \longrightarrow f_2(\varphi(A))$$

to \mathfrak{a}_2 -isomorphisms

$$\mu_A : \varphi'(A) \longrightarrow \varphi(A)$$

with $f_2(\mu_A) = \eta_A$. We can now define $\varphi'(a)$ for $a : A \longrightarrow A'$ in \mathfrak{a}_1 as the unique \mathfrak{a}_2 -map making the following square commute:

$$\begin{array}{ccc} \varphi'(A) & \xrightarrow{\mu_A} & \varphi(A) \\ \varphi'(a) \downarrow & & \downarrow \varphi(a) \\ \varphi'(A') & \xrightarrow{\mu_{A'}} & \varphi(A') \end{array}$$

Clearly, $\varphi' : \mathfrak{a}_1 \longrightarrow \mathfrak{a}_2$ is a functor and $\mu : \varphi' \longrightarrow \varphi$ is a natural isomorphism. If we apply f_2 to the second square and compare the resulting square with the first square, we conclude that

$$f_2 \circ \varphi' = f_1,$$

as we wanted. □

Remark B.2. σ_R need in general not be faithful. Indeed, consider any algebra-deformation $f : A \longrightarrow B$ in which A contains a non-central invertible element ξ with $f(\xi) = 1$. Then $\varphi : A \longrightarrow A : a \longmapsto \xi^{-1}a\xi$ is an algebra-isomorphism different from but naturally isomorphic to 1_A with $f \circ \varphi = f$.

Proposition B.3. *For R in Rng^0/S , σ_R is essentially surjective.*

Proof. Consider a left R -deformation $f : \mathfrak{a} \rightarrow \mathfrak{b}$. Take an inverse equivalence $g : \mathfrak{b} \rightarrow S \otimes_R \mathfrak{a}$ of the canonical equivalence $S \otimes_R f : S \otimes_R \mathfrak{a} \rightarrow \mathfrak{b}$. We construct a category \mathfrak{c} in the following way: the objects of \mathfrak{c} are precisely the objects of \mathfrak{b} . For objects B, B' of \mathfrak{b} , $\mathfrak{c}(B, B')$ is defined to equal $\mathfrak{a}(g(B), g(B'))$ and the composition in \mathfrak{c} is the composition in \mathfrak{a} . There is an obvious functor $\varphi : \mathfrak{c} \rightarrow \mathfrak{a}$ mapping B to $g(B)$ and $a : g(B) \rightarrow g(B')$ to $a : g(B) \rightarrow g(B')$. φ is clearly fully faithful. For A in \mathfrak{a} , $\varphi(f(A)) = g(f(A))$ is isomorphic to A in $S \otimes_R \mathfrak{a}$. But by Proposition A.1, they remain isomorphic in \mathfrak{a} . It follows that φ is essentially surjective and hence an equivalence of categories. Take a natural isomorphism $\eta : 1_{\mathfrak{b}} \rightarrow S \otimes_R f \circ g$. We define a functor $h : \mathfrak{c} \rightarrow \mathfrak{b}$ by putting $h(B) = B$ and mapping $a : g(B) \rightarrow g(B')$ to the unique map $h(a)$ making the following diagram commute:

$$\begin{array}{ccc} B & \xrightarrow{\eta_B} & f(g(B)) \\ h(a) \downarrow & & \downarrow f(a) \\ B' & \xrightarrow{\eta_{B'}} & f(g(B')) \end{array}$$

It follows that the η define a natural isomorphism $h \rightarrow f \circ \varphi$. Finally, we have to show that $S \otimes_R h : S \otimes_R \mathfrak{c} \rightarrow \mathfrak{b}$ is an isomorphism. Since $S \otimes_R h$ is clearly bijective on objects, it suffices that $S \otimes_R h$ or equivalently $S \otimes_R (f \circ \varphi)$ is an equivalence, which is obvious. \square

Theorem B.4. *For R in Rng^0/S , $\text{Sk}(\sigma_R)$ is a bijection.*

Proof. $\text{Sk}(\sigma_R)$ is injective by B.1 and surjective by B.3. \square

REFERENCES

- [1] J. Adámek, H. Herrlich, and G. E. Strecker, *Abstract and concrete categories*, Pure and Applied Mathematics, John Wiley & Sons Inc., New York, 1990, The joy of cats, A Wiley-Interscience Publication.
- [2] M. Artin, A. Grothendieck, and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas, SGA4, Tome 3*, Lecture Notes in Mathematics, vol. 305, Springer Verlag, 1973.
- [3] M. Artin and J. J. Zhang, *Abstract Hilbert schemes*, *Algebr. Represent. Theory* **4** (2001), no. 4, 305–394.
- [4] F. Borceux, *Handbook of categorical algebra. 2*, *Encyclopedia of Mathematics and its Applications*, vol. 51, Cambridge University Press, Cambridge, 1994, Categories and structures.
- [5] N. Bourbaki, *Eléments de mathématique. 22. Première partie: Les structures fondamentales de l'analyse. Livre 1: Théorie des ensembles. Chapitre 4: Structures*, *Actualités Sci. Ind.* no. 1258, Hermann, Paris, 1957.
- [6] M. Gerstenhaber and S. D. Schack, *On the deformation of algebra morphisms and diagrams*, *Trans. Amer. Math. Soc.* **279** (1983), no. 1, 1–50.
- [7] ———, *Algebraic cohomology and deformation theory*, *Deformation theory of algebras and structures and applications (Il Ciocco, 1986)* (Dordrecht), *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, vol. 247, Kluwer Acad. Publ., Dordrecht, 1988, pp. 11–264.
- [8] ———, *The cohomology of presheaves of algebras. I. Presheaves over a partially ordered set*, *Trans. Amer. Math. Soc.* **310** (1988), no. 1, 135–165.
- [9] M. Gerstenhaber, *On the deformation of rings and algebras*, *Ann. of Math. (2)* **79** (1964), 59–103.
- [10] ———, *On the deformation of rings and algebras. II*, *Ann. of Math.* **84** (1966), 1–19.
- [11] A. Grothendieck, *Sur quelques points d'algèbre homologiques*, *Tôhoku Math. J. (2)* **9** (1957), 119–221.
- [12] L. Illusie, *Existence de résolutions globales*, *SGA6, Lecture Notes in Math.*, vol. 225, Springer Verlag, 1971.

- [13] H. Krause, *The spectrum of a module category*, Mem. Amer. Math. Soc. **149** (2001), no. 707, x+125.
- [14] ———, *A Brown representability theorem via coherent functors*, Topology **41** (2002), no. 4, 853–861.
- [15] W. T. Lowen, *A generalization of the Gabriel-Popescu theorem*, to appear.
- [16] ———, *An obstruction theory for objects in abelian categories*, in preparation.
- [17] W. T. Lowen and M. Van den Bergh, *Deformation theory for abelian categories II*, in preparation.
- [18] S. MacLane, *Categories for the working mathematician*, Springer Verlag, Berlin, 1971.
- [19] B. Mitchell, *Rings with several objects*, Advances in Math. **8** (1972), 1–161.
- [20] A. Neeman, *Triangulated categories*, Annals of Mathematics Studies, vol. 148, Princeton University Press, Princeton, NJ, 2001.
- [21] N. Popescu and P. Gabriel, *Caractérisation des catégories abéliennes avec générateurs et limites inductives exactes*, C. R. Acad. Sci. Paris **258** (1964), 4188–4190.
- [22] N. Popescu, *Abelian categories with applications to rings and modules*, Academic Press, London, 1973, London Mathematical Society Monographs, No. 3.
- [23] L. A. Takhtadjan, *Noncommutative homology of quantum tori*, Functional Anal. Appl. **23** (1989), 147–149.

DEPARTEMENT DWIS, VRIJE UNIVERSITEIT BRUSSEL, PLEINLAAN 2, 1050 BRUSSEL, BELGIUM
E-mail address, Wendy T. Lowen: wlowen@vub.ac.be

DEPARTEMENT WNI, LIMBURGS UNIVERSITAIR CENTRUM, UNIVERSITAIRE CAMPUS, BUILDING D, 3590 DIEPENBEEK, BELGIUM
E-mail address, Michel Van den Bergh: vdbergh@luc.ac.be